arXiv:1108.4518v1 [math.AG] 23 Aug 2011

SINGULAR DERIVED CATEGORIES OF $\mathbb{Q}$-FACTORIAL TERMINALIZATIONS AND MAXIMAL MODIFICATION ALGEBRAS

OSAMU IYAMA AND MICHAEL WEMYSS

Abstract. Let X be a Gorenstein normal 3-fold satisfying (ELF) with local rings which are at worst isolated hypersurface (e.g. terminal) singularities. By using the singular derived category $\mathrm{D}_{\mathsf{sg}}(X)$, we give a necessary and sufficient categorical condition for X to be $\mathbb{Q}$-factorial. By passing to the idempotent completion $\overline{\mathrm{D}_{\mathsf{sg}}(X)}$ we obtain a necessary and sufficient condition for X to be complete locally $\mathbb{Q}$-factorial. We then relate this information to maximal modification algebras(=MMAs), introduced in [IW10], by showing that if a modifying algebra Λ is derived equivalent to X as above, then X is $\mathbb{Q}$-factorial if and only if Λ is a MMA. This shows that MMAs categorify $\mathbb{Q}$-factorial terminalizations in the same way that noncommutative crepant resolutions (=NCCRs) categorify 3-fold crepant resolutions. We then apply these results to the case of cA_n singularities, generalizing some results of Burban–Iyama–Keller–Reiten [BIKR] and Dao–Huneke [DH10].

Contents

1. Introduction 1
1.1. Overview 1
1.2. $\mathbb{Q}$-factorial results 3
1.3. General Homological results 4
1.4. cA_n singularities 5
2. Preliminaries 6
2.1. d-CY and d-CY$^-$ 6
2.2. Singular derived categories of algebras 7
2.3. MM and CT modules 8
3. Singular Derived Categories and $\mathbb{Q}$-factorial terminalizations 11
3.1. Preliminaries 11
3.2. Rigid-free, $\mathbb{Q}$-factoriality and MMAs 14
4. cA_n singularities via derived categories 16
4.1. Crepant Modifications of cA_n singularities 16
5. Classification of MM modules on cA_n singularities 19
5.1. Indecomposable modifying modules on cA_n singularities 20
5.2. MM and modifying modules on cA_n singularities 21
5.3. CT modules on cA_n singularities 25
6. Mutation 28
6.1. Some theory 28
6.2. Mutation of modifying generators on cA_n singularities 30
6.3. Geometric corollaries 32
6.4. The mutation graph for maximal modifying modules 32
References 33

1. Introduction

1.1. **Overview.** The purpose of this work, a continuation of [IW10], is to set the foundations of the 'homological MMP', in which the homological and categorical approaches to singularities are used to both (re)prove and also run efficiently certain aspects of the minimal model program (=MMP) in dimension three.

Our first algebraic observation, made and exploited in [IW10], was that homologically, the best setting is that of a maximal modifying module (=MM module), or equivalently a maximal modifying algebra (=MMA). In this situation many strong statements hold, which fail to be true in the more general setting of arbitrary modifying modules. We remarked at the time that this is similar to the geometric phenomenon where the strongest results hold on the $\mathbb{Q}$-factorial terminalizations of $\operatorname{Spec} R$ (which are in the Mori category), as opposed to the more general setting of arbitrary crepant partial resolutions. The first purpose of this paper is to link these two observations.

Before we state our results in detail, it is necessary to first provide a dictionary that links the two fields. Let R be a three-dimensional commutative normal Gorenstein ring (over $\mathbb{C}$), with canonical singularities. Then on the derived level, roughly speaking we always expect correspondences

$$\begin{array}{rcl}
\text{Crepant resolutions of } \operatorname{Spec} R & \longleftrightarrow & \text{CT } R\text{-modules} \\
\rotatebox{90}{$\supseteq$} & & \rotatebox{90}{$\supseteq$} \\
\mathbb{Q}\text{-factorial terminalizations of } \operatorname{Spec} R & \longleftrightarrow & \text{maximal modifying } R\text{-modules} \\
\rotatebox{90}{$\supseteq$} & & \rotatebox{90}{$\supseteq$} \\
\text{Partial crepant resolutions of } \operatorname{Spec} R & \longleftrightarrow & \text{modifying } R\text{-modules}
\end{array}$$

See §1.2 and §2.3 for unexplained terminology. Note that $\mathbb{Q}$-factorial terminalizations of $\operatorname{Spec} R$ always exist [K88, 4.5], only sometimes will they be smooth. Thus the top row is really a special case of the second, since either the top tow exists, in which case the top row and second row are the same, or it does not exist, in which case we study the second row only. Hence, either way, the second row is the natural object of our study. The correspondences take place on the level of the derived categories of the endomorphism rings. More precisely, the endomorphism rings of CT modules give Van den Bergh's non-commutative crepant resolutions (=NCCRs) [IW10, 5.5, 5.12], and all crepant resolutions and all NCCRs are expected to be derived equivalent. Note that this is known in many examples, but even in dimension three it is still not yet known in full generality. Even so, everything in the middle row should be derived equivalent, and furthermore if the top row exists, everything in the top two rows should be derived equivalent. This is known on the algebraic side [IW10, 1.12, 1.15], and is also known on the geometric side [C02]. The content in this paper is linking the two, since this involves characterizing $\mathbb{Q}$-factoriality homologically. Note that in the above picture the bottom row is slightly more subtle than the other two (see for example 6.15), but in the setting of one-dimensional fibres below still behaves well.

Due to the existence of two dimensional fibres, the above are certainly not 1:1 correspondences, however if we restrict ourselves to the cases where the $\mathbb{Q}$-factorial terminalization $Y \to \operatorname{Spec} R$ has only one–dimensional fibres, the picture becomes much more clear. In this case, we expect one–to–one correspondences

$$\begin{array}{rcl}
\text{Crepant resolutions of } \operatorname{Spec} R & \overset{1:1}{\longleftrightarrow} & \text{CT } R\text{-modules} \\
\rotatebox{90}{$\supseteq$} & & \rotatebox{90}{$\supseteq$} \\
\mathbb{Q}\text{-factorial terminalizations of } \operatorname{Spec} R & \overset{1:1}{\longleftrightarrow} & \text{MM } R\text{-modules which are CM} \\
\rotatebox{90}{$\supseteq$} & & \rotatebox{90}{$\supseteq$} \\
\text{Partial crepant resolutions of } \operatorname{Spec} R & \overset{1:1}{\longleftrightarrow} & \text{modifying } R\text{-modules which are CM}
\end{array}$$

where the right hand side is taken up to additive closure. If R is complete local, by restricting to basic modules we can improve on this by taking the right hand side up to isomorphism. We prove parts of this picture for cA_n singularities in §5. Note that again the top row is really a special case of the middle. Also, in the situation of one-dimensional fibres, the existence of a crepant resolution is already known to be equivalent to the existence of a NCCR [V04a] (equivalently, a CT module [IW10, 5.12])

The above picture sets up a correspondence between two fields, allowing us to prove results in one by appealing to simple reasoning in the other. It is our belief that many of the results of the MMP should come out of our categorical picture (along the lines of [Bri02], [BKR]), allowing us to both bypass some of the classifications used in the MMP, and also run some aspects of the MMP in a more efficient manner.

We now describe our results in detail.

1.2. **$\mathbb{Q}$-factorial results.** We begin by studying the $\mathbb{Q}$-factorial property of varieties from the viewpoint of derived categories, then link this to MMAs. These results are independent of the dimension of the fibre, and go some of the way to verifying the second line in the first table.

We denote by $\mathrm{CM}\, R$ the category of CM (=maximal Cohen-Macaulay) R-modules, and by $\underline{\mathrm{CM}}\, R$ its stable category (see §2). As in [Or03, Or09], we denote $\mathrm{D}_{\mathsf{sg}}(X) := \mathrm{D}^{\mathrm{b}}(\mathrm{coh}\, X)/\mathfrak{Perf}(X)$ (see §3.1 for more details). The first key technical result is the following. Note that the idempotent completion part also follows from work of Orlov [Or09], and is well-known to experts [BK11]:

Theorem 1.1. *(=3.7) Suppose that X is a Gorenstein scheme of dimension d satisfying (ELF) (see 3.7 for full details and explanation), with isolated singularities $\{x_1, \dots, x_n\}$. Then there is a fully faithful functor*

$$\mathrm{D}_{\mathsf{sg}}(X) \hookrightarrow \bigoplus_{i=1}^{n} \underline{\mathrm{CM}}\, \mathcal{O}_{X,x_i}$$

which is essentially surjective up to summands. Thus taking the idempotent completion gives an equivalence

$$\overline{\mathrm{D}_{\mathsf{sg}}(X)} \approx \bigoplus_{i=1}^{n} \underline{\mathrm{CM}}\, \widehat{\mathcal{O}}_{X,x_i}.$$

Recall that a normal scheme X is defined to be $\mathbb{Q}$-*factorial* if for every Weil divisor D, there exists $n \in \mathbb{N}$ for which nD is Cartier. If we can always take $n = 1$, we say that X is *locally factorial.* Note that these conditions are Zariski local, i.e. X is $\mathbb{Q}$-factorial if and only if each $\mathcal{O}_{X,x}$ is $\mathbb{Q}$-factorial. It is well–known that $\mathbb{Q}$-factoriality cannot in general be checked complete locally, so we define X to be *complete locally* $\mathbb{Q}$-*factorial* if every completion $\widehat{\mathcal{O}}_{X,x}$ is $\mathbb{Q}$-factorial.

Now if $\mathcal{T}$ is a triangulated category with a suspension functor $[1]$, recall that an object $a \in \mathcal{T}$ is called *rigid* if $\mathrm{Hom}_{\mathcal{T}}(a, a[1]) = 0$. We say that $\mathcal{T}$ is *rigid-free* if every rigid object in $\mathcal{T}$ is isomorphic to the zero object. Even although Zariski locally it is unclear whether the functor in 1.1 is an equivalence, there is still enough information to show the following.

Theorem 1.2. *(=3.9, 3.10) Suppose that X is a normal 3-dimensional Gorenstein scheme over a field k, satisfying (ELF), with isolated singularities $\{x_1, \dots, x_n\}$.*
(1) *If $\mathrm{D}_{\mathsf{sg}}(X)$ is rigid-free, then X is locally factorial.*
(2) *If $\overline{\mathrm{D}_{\mathsf{sg}}(X)}$ is rigid-free, then X is complete locally factorial.*
(3) *If $\mathcal{O}_{X,x_i}$ are hypersurfaces for all $1 \le i \le n$, then the following are equivalent:*
  (a) *X is locally factorial.*
  (b) *X is $\mathbb{Q}$-factorial.*
  (c) *$\mathrm{D}_{\mathsf{sg}}(X)$ is rigid-free.*
  (d) *$\underline{\mathrm{CM}}\, \mathcal{O}_{X,x}$ is rigid-free for all closed points $x \in X$.*
(4) *If $\widehat{\mathcal{O}}_{X,x_i}$ are hypersurfaces for all $1 \le i \le n$, then the following are equivalent:*
  (a) *X is complete locally factorial.*
  (b) *X is complete locally $\mathbb{Q}$-factorial.*
  (c) *$\overline{\mathrm{D}_{\mathsf{sg}}(X)}$ is rigid-free.*
  (d) *$\underline{\mathrm{CM}}\, \widehat{\mathcal{O}}_{X,x}$ is rigid-free for all closed points $x \in X$.*

Rather remarkably, the same homological characterization of rigid-free can be found in the study of modifying and maximal modifying modules (see 2.8 for definitions). Recall that if M is a maximal modifying R-module, we say that $\mathrm{End}_R(M)$ is a *maximal modification algebra (=MMA).*

Proposition 1.3. *(=2.17, 2.3) Suppose that R is normal, Gorenstein, equi-codimensional, 3-dimensional ring, and let M be a modifying R-module. Assume that $\mathrm{End}_R(M)$ has only isolated singularities. Then $\mathrm{End}_R(M)$ is an MMA if and only if the category $\mathrm{D}_{\mathsf{sg}}(\mathrm{End}_R(M))$ is rigid-free.*

Recall that if X is a variety with rational Gorenstein singularities, a $\mathbb{Q}$-factorial terminalization of X is a projective birational morphism $f : Y \to X$ such that Y has only $\mathbb{Q}$-factorial terminal singularities, and further $f^*\omega_X = \omega_Y$. When Y is furthermore smooth, we call $f : Y \to X$ a crepant resolution. Maximal modification algebras were introduced in [IW10] with the aim of generalizing NCCRs to cover the more general situation when crepant resolutions do not exist. We regard the following corollary of 1.2 and 1.3 as one of the main results of this paper.

Theorem 1.4. *Let $Y \to \operatorname{Spec} R$ be a projective birational morphism between normal integral schemes, where R is Gorenstein ring over $\mathbb{C}$ of dimension three, and Y is Gorenstein with only isolated singularities. Suppose that $M \in \operatorname{ref} R$ such that $\operatorname{End}_R(M) \in \operatorname{CM} R$, and further suppose that Y is derived equivalent to $\operatorname{End}_R(M)$. Then*
(1) *If $\operatorname{End}_R(M)$ is a MMA, then Y is locally factorial.*
Further, if Y has locally only isolated hypersurface singularities (e.g. Y has only terminal singularities), then
(2) *Y is $\mathbb{Q}$-factorial $\iff$ $\operatorname{End}_R(M)$ is a MMA.*

Thus MMAs categorify 3-fold $\mathbb{Q}$-factorial terminalizations in the same way that NCCRs categorify 3-fold crepant resolutions. Hence, following Bondal–Orlov and Van den Bergh, using the above results we conjecture the following:

Conjecture 1.5. Suppose that R is a normal Gorenstein 3-fold over $\mathbb{C}$, with only canonical (equivalently rational, since R is Gorenstein [R87, (3.8)]) singularities. Then all $\mathbb{Q}$-factorial terminalizations of R and all maximal modification algebras are derived equivalent.

We point out that, if true, then Conjecture 1.5 implies that all crepant resolutions of R (both commutative and non-commutative) are derived equivalent, settling a conjecture of Van den Bergh [V04b, 4.6]. We remark that 1.5 has some evidence:

Theorem 1.6. *Suppose that R is a normal Gorenstein 3-fold over $\mathbb{C}$ whose $\mathbb{Q}$-factorial terminalizations $Y \to \operatorname{Spec} R$ have only one dimensional fibres (e.g. R is a terminal Gorenstein singularity, or a cA_n singularity in §4). Then Conjecture 1.5 is true.*

Proof. The $\mathbb{Q}$-factorial terminalizations are connected by flops, so they are derived equivalent by [C02]. By [V04a] there is a derived equivalence between Y and some algebra $\operatorname{End}_R(M)$ with $\operatorname{End}_R(M) \in \operatorname{CM} R$. Thus by 1.4 $\operatorname{End}_R(M)$ is a MMA. We already know that all MMAs are derived equivalent [IW10, 4.9]. □

As it stands, the proof of 1.6 heavily uses the MMP. We believe that it should be possible to prove 1.6 without using the MMP, instead using the categorical techniques developed in this paper.

1.3. General Homological results.

In the homological MMP, the technique of (categorical) mutation developed in [IW10, §6] is the operation corresponding to flop. However, mutation can be applied to much more general situations than just flopping contractions, and it is this flexibility that also allows us to generate derived equivalences in both the highly singular and the more general contexts.

In the theory of mutation developed in [IW10, §6], the main setting was dimension three, in which the behaviour depends heavily on the dimension (as a vector space) of a certain factor algebra Λ_J. Our new result is the following, which in fact works in full generality. We refer the reader to §6 for the definition of the mutation operator ν_J, and all other unexplained terminology.

Theorem 1.7. *(=6.3, 6.6) Let R be a complete local normal d-CY^- commutative ring, $M := R \oplus (\oplus_{i\in I} M_i)$ be a basic modifying R-module and choose any $\emptyset \neq J \subseteq I$. Then $\nu_J(M)$ is a modifying module, and further $\operatorname{End}_R(M)$ is derived equivalent to $\operatorname{End}_R(\nu_J(M))$.*

This allows us to mutate at will, without having the rather delicate task of determining whether $\dim_k \Lambda_J < \infty$. We calculate the mutations (in this level of generality) for cA_n singularities in §6.

1.4. cA_n **singularities.** Suppose that $f_1,\ldots,f_n \in \mathfrak{m} := (x,y) \subseteq k[[x,y]]$ are (not necessarily distinct) irreducible polynomials, and let $R := k[[u,v,x,y]]/(uv - f_1 \ldots f_n)$. For each $\omega \in \mathfrak{S}_n$, define

$$T^\omega := R \oplus (u, f_{\omega(1)}) \oplus (u, f_{\omega(1)} f_{\omega(2)}) \oplus \ldots \oplus (u, f_{\omega(1)} \ldots f_{\omega(n-1)}).$$

Our first result is the following generalization of [BIKR, 1.5] and [DH10, 4.2], which follows more or less immediately from 1.4(2).

Theorem 1.8. *(=4.4) Let $k = \mathbb{C}$ and suppose that $f_1,\ldots,f_n \in \mathfrak{m} := (x,y) \subseteq \mathbb{C}[[x,y]]$ are irreducible polynomials. Let $R = \mathbb{C}[[u,v,x,y]]/(uv - f_1 \ldots f_n)$, then*
(1) Each T^ω ($\omega \in \mathfrak{S}_n$) is a MM R-module generator. The corresponding endomorphism rings all have isolated singularities.
(2) T^ω are CT modules $\iff$ $f_i \notin \mathfrak{m}^2$ for all $1 \leq i \leq n$.

Using commutative algebra, we then give an independent (and strengthened) proof of 1.8, which does not use the explicit form of the $\mathbb{Q}$-factorial terminalizations of Spec R, and also is valid over an arbitrary algebraically closed field $\Bbbk$. This is important, since in many other examples the explicit forms of $\mathbb{Q}$-factorial terminalizations are not known. The strengthened version is as follows:

Theorem 1.9. *(=5.17, 5.19) Suppose that $f_1,\ldots,f_n \in \mathfrak{m} := (x,y) \subseteq \Bbbk[[x,y]]$ are irreducible polynomials. Let $R = \Bbbk[[u,v,x,y]]/(uv - f_1 \ldots f_n)$, then*
(1) The MM R-module generators are precisely T^ω, where $\omega \in \mathfrak{S}_n$.
(2) R has a CT module $\iff$ $f_i \notin \mathfrak{m}^2$ for all $1 \leq i \leq n$. In this case, the CT R-modules are precisely T^ω, where $\omega \in \mathfrak{S}_n$.

There were no examples of MMAs given in [IW10] (except in §6, which is actually an NCCR), so to make MMAs more concrete we give (in 5.21) the explicit quiver of the endomorphism ring of T^ω.

Keeping R as above, we then produce many examples of derived equivalences between the partial crepant resolutions of Spec R, which in general have canonical singularities. The partial crepant resolutions of Spec R have a certain number of curves, and on these curves singularities of the form $uv = f_I$ (see 4.6). We denote the partial resolutions combinatorially in terms of flags $\mathcal{F}$, and denote the corresponding spaces by $X^{\mathcal{F}}$ (see §4.1).

Theorem 1.10. *(=6.16) $X^{\mathcal{F}}$ and $X^{\mathcal{G}}$ are derived equivalent if they have the same number of curves above the origin of* Spec R*, and the singularities of $X^{\mathcal{F}}$ can be permuted to the singularities of $X^{\mathcal{G}}$.*

In fact, 1.10 comes very easily from a simple calculation which determines the mutations of a given modifying module:

Theorem 1.11. *(=6.13) Fix a flag $\mathcal{F} = (I_1,\ldots,I_m)$, and associate to $\mathcal{F}$ the module $T^{\mathcal{F}}$ and the combinatorial picture $\mathcal{P}(\mathcal{F})$. Choose $\emptyset \neq J \subseteq \{1,\ldots,m\}$, then $\nu_J(T^{\mathcal{F}})$ is the module corresponding to the J-reflection of $\mathcal{P}(\mathcal{F})$.*

See §6 for full details. In particular, since we prove in 6.17 that the mutation graph of MM modules is connected, 1.10 gives the following alternative proof of [C02] in the case of complete local cA_n singularities, which does not involve passing to dimension four:

Corollary 1.12. *Let $R = \mathbb{C}[[u,v,x,y]]/(uv - f_1 \ldots f_n)$, then all $\mathbb{Q}$-factorial terminalizations of* Spec R *are derived equivalent.*

We remark that although the above is in the complete local setting, this is mainly for our own convenience, since it simplifies calculations. We believe that once the proof is made fully homological, 1.12 will generalize.

Conventions. Throughout R will always denote a *commutative* ring. All modules will be left modules, so for a ring A we denote mod A to be the category of finitely generated left A-modules, and Mod A will denote the category of all left A-modules. Throughout when composing maps fg will mean f then g, similarly for quivers ab will mean a then b. Note that with this convention $\mathrm{Hom}_R(M,X)$ is a $\mathrm{End}_R(M)$-module and $\mathrm{Hom}_R(X,M)$

is a $\mathrm{End}_R(M)^{\mathrm{op}}$-module. For $M \in \mathrm{mod}\, A$ we denote $\mathrm{add}\, M$ to be the full subcategory consisting of summands of finite direct sums of copies of M, and we denote $\mathrm{proj}\, A := \mathrm{add}\, A$ to be the category of finitely generated projective A-modules. We say that $M \in \mathrm{mod}\, R$ is a *generator* if $R \in \mathrm{add}\, M$.

If X is a scheme, $\mathcal{O}_{X,x}$ will denote the localization of the structure sheaf at the closed point $x \in X$. We will denote by $\widehat{\mathcal{O}}_{X,x}$ the completion of $\mathcal{O}_{X,x}$ at the unique maximal ideal. For us, *locally* will always mean Zariski locally, that is if we say that R has locally only isolated hypersurface singularities, we mean that each $R_{\mathfrak{m}}$ is an isolated hypersurface singularity (or is smooth). When we want to discuss the completion, we will always refer to this as *complete locally.*

Rings and schemes will *not* be assumed to be finite type over a field k, unless specified. When we say 'over k' we mean 'of finite type over k'. Throughout, k will denote an arbitrary field, and K will denote a field of characteristic zero. Bold letters will denote algebraically closed fields, so $\Bbbk$ will denote an arbitrary algebraically closed field, and $\mathbb{K}$ will denote an algebraically closed field of characteristic zero.

Acknowledgements. The authors would like to thank Vanya Cheltsov, Hailong Dao, Anne-Sophie Kaloghiros, Raphael Rouquier and Ed Segal for many invaluable discussions regarding this work.

2. PRELIMINARIES

For a commutative noetherian local ring $(R, \mathfrak{m})$ and $M \in \mathrm{mod}\, R$, recall that the *depth* of M is defined to be $\mathrm{depth}_R\, M := \inf\{i \geq 0 : \mathrm{Ext}^i_R(R/\mathfrak{m}, M) \neq 0\}$. We say that $M \in \mathrm{mod}\, R$ is *maximal Cohen-Macaulay* (or simply, *CM*) if $\mathrm{depth}_R\, M = \dim R$. Now let R be a (not necessarily local) commutative noetherian ring. We say that $M \in \mathrm{mod}\, R$ is *CM* if $M_{\mathfrak{p}}$ is CM for all prime ideals $\mathfrak{p}$ in R, and we say that R is a *CM ring* if R is a CM R-module. Denoting $(-)^* := \mathrm{Hom}_R(-, R) : \mathrm{mod}\, R \to \mathrm{mod}\, R$, we say that $X \in \mathrm{mod}\, R$ is *reflexive* if the natural map $X \to X^{**}$ is an isomorphism.

We denote $\mathrm{ref}\, R$ to be the category of reflexive R-modules, and we denote $\mathrm{CM}\, R$ to be the category of CM R-modules.

2.1. d-CY and d-CY$^-$.

When dealing with singularities, throughout this paper we use the language of d-CY$^-$ algebras, introduced in [IR08, §3]. Although technical, it provides the common language that links the commutative rings and the various noncommutative endomorphism rings that we will consider.

To do this, let R be a commutative noetherian ring with $\dim R = d$ and let Λ be a module-finite R-algebra. For any $X \in \mathrm{mod}\, \Lambda$, denote by $E(X)$ the injective hull of X, and put $E := \oplus_{\mathfrak{m} \in \mathrm{Max}\, R} E(R/\mathfrak{m})$. This gives rise to Matlis duality $D := \mathrm{Hom}_R(-, E)$. Note that if R is over $\Bbbk$, then $D = \mathrm{Hom}_{\Bbbk}(-, \Bbbk)$ by [O76, 1.1, 1.2]. We let $\mathrm{D}^{\mathrm{b}}_{\mathsf{fl}}(\Lambda)$ denote all bounded complexes with finite length cohomology.

Definition 2.1. *For $n \in \mathbb{Z}$, we call Λ n-CY if there is a functorial isomorphism*

$$\mathrm{Hom}_{\mathrm{D}(\mathrm{Mod}\,\Lambda)}(X, Y[n]) \cong D\, \mathrm{Hom}_{\mathrm{D}(\mathrm{Mod}\,\Lambda)}(Y, X)$$

for all $X, Y \in \mathrm{D}^{\mathrm{b}}_{\mathsf{fl}}(\Lambda)$. Similarly we call Λ n-CY$^-$ if the above functorial isomorphism holds for all $X \in \mathrm{D}^{\mathrm{b}}_{\mathsf{fl}}(\Lambda)$ and $Y \in \mathrm{K}^{\mathrm{b}}(\mathrm{proj}\, \Lambda)$.

We know [IR08, 3.1(7)] that Λ is n-CY if and only if it is n-CY$^-$ and $\mathrm{gl.dim}\, \Lambda < \infty$. Also, by [IR08, 3.10], if R is a commutative noetherian ring and $d \in \mathbb{N}$, Then R is d-CY$^-$ if and only if R is Gorenstein and equi-codimensional, with $\dim R = d$.

Definition 2.2. *Let Λ be a module-finite R-algebra where R is an arbitrary commutative ring. We call Λ a* symmetric R-algebra *if $\mathrm{Hom}_R(\Lambda, R) \cong \Lambda$ as Λ-bimodules. We call Λ a* locally symmetric R-algebra *if $\Lambda_{\mathfrak{p}}$ is a symmetric $R_{\mathfrak{p}}$-algebra for all $\mathfrak{p} \in \mathrm{Spec}\, R$.*

We also say that Λ is an *R-order* if $\Lambda \in \mathrm{CM}\, R$. Now if $\Lambda := \mathrm{End}_R(M)$, where R is normal d-CY$^-$ and $M \in \mathrm{ref}\, R$, then Λ is automatically symmetric by [IR08, 2.4(3)]. If further $\Lambda \in \mathrm{CM}\, R$, then certainly Λ is a locally symmetric R-order, hence [IR08, 3.2(2)] implies that $\Lambda_{\mathfrak{m}}$ is d-CY$^-$ for all $\mathfrak{m} \in \mathrm{Max}\, R$, hence Λ is d-CY$^-$ [IR08, 3.3(1)].

2.2. **Singular derived categories of algebras.** Suppose that R is a d-CY^{-} commutative noetherian ring, and let Λ be a d-CY^{-} module-finite R-algebra. We define $X \in \operatorname{mod}\Lambda$ to be a CM Λ-module if $\operatorname{Ext}^i_\Lambda(X,\Lambda)=0$ for all $i \geq 1$. The proof of [IR08, 3.4(5)(i)] shows that $\operatorname{Ext}^i_\Lambda(X,\Lambda)=0 \iff \operatorname{Ext}^i_R(X,R)=0$. Hence X is a CM Λ-module if and only if it is CM as an R-module. We denote by $\operatorname{CM}\Lambda$ the category of CM Λ-modules, and we denote $\underline{\operatorname{CM}}\Lambda$ to be the stable category, where we factor out by those morphisms which factor through projective Λ-modules.

When Λ is a noetherian ring, we define $\mathrm{D}_{\mathsf{sg}}(\Lambda) := \mathrm{D}^{\mathrm{b}}(\operatorname{mod}\Lambda)/\mathrm{K}^{\mathrm{b}}(\operatorname{proj}\Lambda)$. The following is a now standard theorem of Buchweitz, and is very well known:

Theorem 2.3. *Suppose that Λ is d-CY^{-}, then*

$$\mathrm{D}_{\mathsf{sg}}(\Lambda) \approx \underline{\operatorname{CM}}\,\Lambda.$$

Proof. Since $\Lambda = \operatorname{End}_R(M)$ is d-CY^{-}, by [IR08, 3.1(6)] $\operatorname{inj.dim}_\Lambda \Lambda < \infty$. Similarly, since Λ^{op} is d-CY^{-} [IR08, 3.1(2)], $\operatorname{inj.dim}_{\Lambda^{\mathrm{op}}} \Lambda^{\mathrm{op}} < \infty$. Hence by definition Λ is strongly Gorenstein in the sense of [Bu86, above 4.1.1], and so the result is a special case of [Bu86, 4.4.1(2)]. ☐

Recall [Aus84] that we say an R-algbera Λ has *isolated singularities* if $\operatorname{gl.dim}\Lambda_{\mathfrak{p}} < \infty$ for all non-maximal primes $\mathfrak{p}$ of R.

Proposition 2.4. *Suppose that Λ is d-CY^{-}. Then Λ has isolated singularities if and only if all Hom-spaces in $\mathrm{D}_{\mathsf{sg}}(\Lambda)$ are finite length R-modules.*

Proof. This is well-known (e.g. [Yos90, 3.3], [Aus84]), but for convenience we give the proof.
($\Rightarrow$) By 2.3 all Hom-spaces in $\mathrm{D}_{\mathsf{sg}}(\Lambda)$ are of the form $\operatorname{Hom}_{\underline{\operatorname{CM}}\,\Lambda}(X,Y)$ for some $X,Y \in \operatorname{CM}\Lambda$. Since $\operatorname{Hom}_{\underline{\operatorname{CM}}\,\Lambda}(X,Y) \cong \operatorname{Ext}^1_\Lambda(\Omega^{-1}X,Y)$ and $\Omega^{-1}X \in \operatorname{CM}\Lambda$, the result is clear.
($\Leftarrow$) Let $\mathfrak{p}$ be a prime with $\operatorname{ht}\mathfrak{p} < d$ and let $X \in \operatorname{mod}\Lambda_{\mathfrak{p}}$. Certainly we can find $Y \in \operatorname{mod}\Lambda$ with $Y_{\mathfrak{p}} \cong X$ and so consider a projective resolution

$$0 \to K \to P_{d-1} \to \ldots \to P_1 \to P_0 \to Y \to 0$$

in $\operatorname{mod}\Lambda$. Since $\Lambda \in \operatorname{CM}R$, by localizing and using the depth lemma we see that $K \in \operatorname{CM}R$, i.e. $K \in \operatorname{CM}\Lambda$. Consider

$$0 \to \Omega K \to P_d \to K \to 0, \tag{2.A}$$

then since $\operatorname{Ext}^1_\Lambda(K,\Omega K) \cong \operatorname{Hom}_{\underline{\operatorname{CM}}\,\Lambda}(\Omega K,\Omega K)$ has finite length, it is supported only on maximal ideals. Hence (2.A) splits under localization to $\mathfrak{p}$ and so $K_{\mathfrak{p}}$ is free. Thus $\operatorname{proj.dim}_{\Lambda_{\mathfrak{p}}} X < d$, as required. ☐

The following is based on some Danish handwritten notes of Hans-Bjørn Foxby.

Theorem 2.5. *Let R be a commutative noetherian ring (not necessarily of finite Krull dimension), let Λ be a module-finite R-algebra. and let $M \in \operatorname{mod}\Lambda$. If $\operatorname{proj.dim}_{\Lambda_{\mathfrak{p}}}(M_{\mathfrak{p}}) < \infty$ for all $\mathfrak{p} \in \operatorname{Spec}R$, then $\operatorname{proj.dim}_\Lambda M < \infty$.*

Proof. Take a finitely generated projective resolution of M,

$$\ldots \to P_1 \xrightarrow{d_1} P_0 \xrightarrow{d_0} M \to 0,$$

and set $K_0 = M = \operatorname{Im} d_0$ and $K_i = \operatorname{Im} d_i = \operatorname{Ker} d_{i-1}$ for $i > 0$. Now

$$\operatorname{proj.dim}_\Lambda(M) \leq n \iff \operatorname{Ext}^{n+1}_\Lambda(M,K_{n+1}) = 0,$$

and the right hand side is equivalent to showing that $\operatorname{Supp}_R \operatorname{Ext}^{n+1}_\Lambda(M,K_{n+1})$ is empty. Now since $\operatorname{Ext}^{n+1}_\Lambda(M,K_{n+1})$ is a finitely generated R-module

$$\begin{aligned}\operatorname{proj.dim}_{\Lambda_{\mathfrak{p}}}(M_{\mathfrak{p}}) > n &\iff \operatorname{Ext}^{n+1}_\Lambda(M,K_{n+1}))_{\mathfrak{p}} = \operatorname{Ext}^{n+1}_{\Lambda_{\mathfrak{p}}}(M_{\mathfrak{p}},(K_{n+1})_{\mathfrak{p}}) \neq 0 \\ &\iff \mathfrak{p} \in \operatorname{Supp}_R \operatorname{Ext}^{n+1}_\Lambda(M,K_{n+1}).\end{aligned}$$

But again $\operatorname{Ext}^{n+1}_\Lambda(M,K_{n+1})$ is a finitely generated R-module, hence

$$\operatorname{Supp}_R \operatorname{Ext}^{n+1}_\Lambda(M,K_{n+1}) = V(\operatorname{Ann}_R(\operatorname{Ext}^{n+1}_\Lambda(M,K_{n+1}))).$$

For ease of notation set $I_n = \mathrm{Ann}_R(\mathrm{Ext}^{n+1}_\Lambda(M, K_{n+1}))$, then

$$\mathrm{proj.dim}_{\Lambda_\mathfrak{p}}(M_\mathfrak{p}) > n \iff \mathfrak{p} \in V(I_n). \tag{2.B}$$

In particular,

$$V(I_0) \supseteq V(I_1) \supseteq V(I_2) \supseteq \dots,$$

from which it follows [H77, II.2.1] that

$$\mathrm{Rad}(I_0) \subseteq \mathrm{Rad}(I_1) \subseteq \mathrm{Rad}(I_2) \subseteq \dots \tag{2.C}$$

Let I be the union of the chain in (2.C); its an ideal since its the union of a chain of ideals. It is clear that $V(I) = V(\bigcup_{n=0}^\infty \mathrm{Rad}(I_n)) = \bigcap_{n=0}^\infty V(\mathrm{Rad}(I_n)) = \bigcap_{n=0}^\infty V(I_n)$. Now if $\mathfrak{p} \in V(I)$, then $\mathfrak{p} \in V(I_n)$ for all n, hence by (2.B) $\mathrm{proj.dim}_{\Lambda_\mathfrak{p}}(M_\mathfrak{p}) = \infty$. But this is a contradiction, hence $V(I)$ must be empty, and thus $I = R$. Consequently $I_t = R$ for some t, and so $\mathrm{Ext}^{t+1}_\Lambda(M, K_{t+1}) = 0$. ☐

Corollary 2.6. *Suppose that Λ is a module-finite R-algebra. Then*
(1) For every $\mathfrak{m} \in \mathrm{Max}\, R$, $\Lambda_\mathfrak{m}$ has only finitely many simple modules.
(2) Suppose that $K \in \mathrm{mod}\,\Lambda$ such that for all $Y \in \mathrm{mod}\,\Lambda$, $\mathrm{Ext}^j_\Lambda(K, Y) = 0$ for $j >> 0$. Then $\mathrm{proj.dim}_\Lambda K < \infty$.

Proof. (1) $\Lambda_\mathfrak{m}$ is a module-finite $R_\mathfrak{m}$-algebra. There are only finitely many simple $\Lambda_\mathfrak{m}$-modules since they are annihilated by $\mathfrak{m}$ and hence they are modules over the finite dimensional $R/\mathfrak{m}$-algebra $\Lambda/\mathfrak{m}\Lambda$, which has only finitely many simples.
(2) We know that $\mathrm{mod}\,\Lambda \to \mathrm{mod}\,\Lambda_\mathfrak{m}$ is essentially surjective, so let $Y \in \mathrm{mod}\,\Lambda$ be such that $Y_\mathfrak{m}$ is the sum of the (finite number of) simple $\Lambda_\mathfrak{m}$-modules. The assumptions imply that $\mathrm{Ext}^j_{\Lambda_\mathfrak{m}}(K_\mathfrak{m}, Y_\mathfrak{m}) = 0$ for high enough j, hence $\mathrm{proj.dim}_{\Lambda_\mathfrak{m}} K_\mathfrak{m} < \infty$. This is true for all $\mathfrak{m} \in \mathrm{Max}\, R$, hence $\mathrm{proj.dim}_{\Lambda_\mathfrak{p}} K_\mathfrak{p} < \infty$ for all $\mathfrak{p} \in \mathrm{Spec}\, R$. By 2.5, it follows that $\mathrm{proj.dim}_\Lambda K < \infty$. ☐

Now recall that $X \in \mathfrak{T}$ is called *homologically finite* if for all $y \in \mathfrak{T}$, $\mathrm{Hom}_\mathfrak{T}(x, y[i]) = 0$ for all but finitely many i.

Lemma 2.7. *Suppose that Λ is a module-finite R-algebra. Then the homologically finite complexes in $\mathrm{D^b}(\mathrm{mod}\,\Lambda)$ are precisely $\mathrm{K^b}(\mathrm{proj}\,\Lambda)$.*

Proof. Certainly every member of $\mathrm{K^b}(\mathrm{proj}\,\Lambda)$ is homologically finite, we just need the converse. If X is homologically finite, replace X by its projective resolution (P, d). The truncation of P, namely $\tau_{\geq i}P$, is quasi-isomorphic to X for large enough i. Fix such an i. Thus, if we can show that $\ker(d_i)$ has finite projective dimension, certainly it follows that X belongs to $\mathrm{K^b}(\mathrm{proj}\,\Lambda)$.

Now by the usual short exact sequence given by truncation, there is a triangle

$$Q \to X \to \ker(d_i)[i] \to Q[1]$$

where $Q \in \mathrm{K^b}(\mathrm{proj}\,\Lambda)$. Since the first two are homologically finite, so is $\ker(d_i)$. In particular, for all $Y \in \mathrm{mod}\,\Lambda$, we have that $\mathrm{Ext}^j_\Lambda(\ker(d_i), Y) = 0$ for large enough j. By 2.6(2), $\ker(d_i)$ has finite projective dimension, as required. ☐

2.3. MM and CT modules.

We briefly recap the definitions of modifying, maximal modifying, and CT modules, then give some basic lemmas.

Definition 2.8. [IW10] *Let R be a d-CY$^-$ ring. Then*
(1) $X \in \mathrm{ref}\, R$ is called a modifying module *if $\mathrm{End}_R(M) \in \mathrm{CM}\, R$.*
(2) We say $X \in \mathrm{ref}\, R$ is a maximal modifying (=MM) module *if it is modifying and further if $X \oplus Y$ is modifying for $Y \in \mathrm{ref}\, R$, then $Y \in \mathrm{add}\, X$. Equivalently,*

$$\mathrm{add}\, M = \{X \in \mathrm{ref}\, R \mid \mathrm{End}_R(M \oplus X) \in \mathrm{CM}\, R\}.$$

(3) We call $M \in \mathrm{CM}\, R$ a CT module *if*

$$\mathrm{add}\, M = \{X \in \mathrm{CM}\, R \mid \mathrm{Hom}_R(M, X) \in \mathrm{CM}\, R\}.$$

The following three results will be used extensively. As in [IW10], if $X \in \mathrm{mod}\, R$, we denote $\mathsf{fl}\, X$ to be the largest finite length R-submodule of X.

Lemma 2.9. [IW10, 2.6] *Suppose that R is a 3-CY$^-$ ring, and let $M \in \operatorname{ref} R$.*
(1) *If M is modifying, then $\operatorname{fl}\operatorname{Ext}^1_R(M,M)=0$. The converse holds if $M \in \operatorname{CM} R$.*
(2) *Assume that R is an isolated singularity. If M is modifying, then it is rigid. The converse holds if $M \in \operatorname{CM} R$.*

Lemma 2.10. [IW10, 2.18(2)] *Suppose that R is a d-CY$^-$ normal domain, and $M \in \operatorname{ref} R$. Then M is modifying if and only if $\operatorname{End}_R(M)$ is d-CY$^-$.*

Proposition 2.11. [IW10, 5.14(2)] *Let R be a 3-CY$^-$ normal domain, and assume that R has a CT module. Then the MM modules which are CM (equivalently, the MM generators) are precisely the CT modules.*

It is not clear from the definition, but it is nevertheless true [IW10, 5.5], that the property CT can be checked complete locally, that is M is a CT R-module if and only if $\widehat{M}_{\mathfrak{m}}$ is a CT $\widehat{R}_{\mathfrak{m}}$-module for all $\mathfrak{m} \in \operatorname{Max} R$. In §3, this corresponds to the fact that a scheme X is non-singular if and only if complete locally it is non-singular.

We do not know whether the property of being MM can be checked locally, but certainly it cannot be checked complete locally. One direction is true, namely if $\widehat{M}_{\mathfrak{m}}$ is a MM $\widehat{R}_{\mathfrak{m}}$-module for all $\mathfrak{m} \in \operatorname{Max} R$, then M is a MM R-module. However, it is precisely the failure of the converse that is the algebraic phenomenon corresponding to the difference between $\mathbb{Q}$-factorial and complete locally $\mathbb{Q}$-factorial singularities in §3.

Example 2.12. Consider the element $f = x^2+x^3+y^2-uv$ in the ring $\mathbb{C}[u,v,x,y]$, and set $R := \mathbb{C}[u,v,x,y]/(f)$. Then R is a maximal modifying R-module, but $\widehat{R}_{\mathfrak{m}}$ is not a maximal modifying $\widehat{R}_{\mathfrak{m}}$-module, where $\mathfrak{m}=(u,v,x,y)$.

Proof. Suppose M is a modifying R-module. We show that M is projective. Since R has an isolated singularity only at $\mathfrak{m}$, we know that $\operatorname{Ext}^1_R(M,M)=0$, and M is locally free on the punctured spectrum. Thus we just need to prove that $M_{\mathfrak{m}}$ is a free $R_{\mathfrak{m}}$-module. Certainly $\operatorname{Ext}^1_{R_{\mathfrak{m}}}(M_{\mathfrak{m}},M_{\mathfrak{m}})=0$ and so $M_{\mathfrak{m}}$ is a modifying $R_{\mathfrak{m}}$-module. But $R_{\mathfrak{m}}$ is a local Gorenstein factorial domain (since $\sqrt{x+1}$ does not exist Zariski locally), hence by 2.14 $M_{\mathfrak{m}}$ is free.

For the last statement, since $\sqrt{x+1}$ exists (and is a unit) in the completion, $\widehat{R}_{\mathfrak{m}}$ is isomorphic to $\mathbb{C}[[u,v,x,y]]/(uv-xy)$, for which $\widehat{R}_{\mathfrak{m}} \oplus (u,x)$ is a modifying module. ☐

We record here the following easy lemma, which shows the first link between modifying objects and factoriality. Recall that $\underline{\operatorname{CM}} R$ denotes the stable category of CM modules, where we factor out by those morphisms which factor through projective R-modules.

Lemma 2.13. *Suppose R is 3-CY$^-$ normal domain, with isolated singularities. Consider the statements*
(1) *R is a MM module.*
(2) *Every modifying R-module is projective.*
(3) *$\underline{\operatorname{CM}} R$ is rigid-free.*
(4) *R is locally factorial.*
Then we have (1) ⇔ (2) ⇔ (3) ⇒ (4).

Proof. (1)⇒(2) Let $M \in \operatorname{ref} R$ be any modifying module, then since R is maximal we may find a 2-term R-approximation of M [IW10, 4.6]

$$0 \to F_1 \to F_0 \xrightarrow{f} M$$

with $F_i \in \operatorname{add} R = \operatorname{proj}(R)$. Since this is an R-approximation, necessarily f is surjective. Thus $\operatorname{proj.dim}_R M \le 1$. By [AG60, 4.10], if $\operatorname{proj.dim}_R M = 1$ then $\operatorname{Ext}^1_R(M,M) \neq 0$. However, since R is isolated, $\operatorname{Ext}^1_R(M,M) = 0$ by 2.9(2). Hence M is projective, as required.
(2)⇒(3) is clear, since $\operatorname{Hom}_{\underline{\operatorname{CM}} R}(M,M[1]) \cong \operatorname{Ext}^1_R(M,M)$.
(3)⇒(1) Suppose that $\operatorname{End}_R(R\oplus X) \in \operatorname{CM} R$ for some $X \in \operatorname{ref} R$. Necessarily $X \in \operatorname{CM} R$ with X modifying. Hence X is zero in $\underline{\operatorname{CM}} R$, i.e. $X \in \operatorname{add} R$.
(2)⇒(4) Since R is a normal domain, all members of the class group are modifying R-modules. Hence every member of the class group is projective, so the result follows. ☐

In future sections we will often use the following result from commutative algebra:

Theorem 2.14. [Dao1, 3.1(1)] *Let $R = S/(f)$ where S is a regular local ring of dimension four, containing a field k, and f is a regular element in S. If R is a $\mathbb{Q}$-factorial isolated singularity, then the free modules are the only modifying modules.*

Proof. This is the equi-characteristic version of [Dao1, 3.1(1)], but since *loc. cit.* is in somewhat different language than used here, so we sketch Dao's proof. Let $N \in \operatorname{ref} R$ such that $\operatorname{End}_R(N) \in \operatorname{CM} R$. Since R is isolated, $\operatorname{Ext}^1_R(N,N) = 0$ by 2.9(2). On the other hand, by [Dao2, 4.1], N is a Tor-rigid R-module. These two facts imply, via a result of Jothilingham [Jo75, Main Theorem] (see also [Dao1, 2.4]), that N is free R-module. $\square$

Corollary 2.15. *Let R be a 3-CY^- normal domain over k, which locally has only isolated hypersurface singularities. Then the following are equivalent:*
(1) R is a MM module.
(2) Every modifying R-module is projective.
(3) $\underline{\operatorname{CM}} R$ is rigid-free.
(4) R is locally factorial.

Proof. (1)⇔(2)⇔(3)⇒(4) is 2.13.
(4)⇒(1), suppose that $M \in \operatorname{ref} R$ with $\operatorname{End}_R(R \oplus M) \in \operatorname{CM} R$. Then $\operatorname{End}_{R_\mathfrak{m}}(R_\mathfrak{m} \oplus M_\mathfrak{m}) \in \operatorname{CM} R_\mathfrak{m}$ with $R_\mathfrak{m}$ satisfying the assumptions of 2.14, hence $M_\mathfrak{m}$ is a free $R_\mathfrak{m}$-module. Thus $M_\mathfrak{m} \in \operatorname{add} R_\mathfrak{m}$ for all $\mathfrak{m} \in \operatorname{Max} R$, so $M \in \operatorname{add} R$ [IW10, 4.11]. $\square$

Remark 2.16. The corollary is false if we remove the isolated singularities assumption, since then it is not necessarily true that (4) implies (1). A concrete example is $R = \mathbb{C}[x,y,z,t]/(x^2+y^3+z^5)$. Also, note that in the isolated singularity case it is a little unclear if the hypersurface assumption is strictly necessary; indeed Dao conjectures his result still holds in the case of complete intersections [Dao3, §4]. By itself, the statement in 2.14 does not refer to Tor–rigidity, so may even be true in full generality. We remark that removing the hypersurface assumption in 2.15 would allow us to bypass the use of the classification of terminal singularities later.

We note that the maximality of a given modifying module can be determined categorically, generalizing 2.13.

Proposition 2.17. *Suppose that R is normal 3-CY^-, and let M be a modifying R-module. Set $\Lambda_M := \operatorname{End}_R(M)$. Then*
(1) M is a MM R-module $\iff$ $\operatorname{fl}\operatorname{Hom}_{\underline{\operatorname{CM}}\Lambda_M}(Y, Y[1]) \neq 0$ for all $0 \neq Y \in \underline{\operatorname{CM}}\Lambda_M$.
If further Λ_M has isolated singularities, then
(2) M is a MM R-module $\iff$ $\operatorname{Hom}_{\underline{\operatorname{CM}}\Lambda_M}(Y, Y[1]) \neq 0$ for all $0 \neq Y \in \underline{\operatorname{CM}}\Lambda_M$.
Hence if Λ_M has isolated singularities, then M is maximal modifying if and only if the category $\underline{\operatorname{CM}}\Lambda_M$ is rigid-free.

Proof. (1) By reflexive equivalence, there exists $X \in \operatorname{ref} R$ such that $\operatorname{End}_R(M \oplus X) \in \operatorname{CM} R$ if and only if there exists $X \in \operatorname{ref} R$ such that $\operatorname{End}_{\Lambda_M}(\Lambda_M \oplus \operatorname{Hom}_R(M,X)) \in \operatorname{CM} R$. Now $\Lambda_M \in \operatorname{CM} R$ by assumption, further

$$\operatorname{Hom}_{\Lambda_M}(\Lambda_M, \operatorname{Hom}_R(M,X)) \in \operatorname{CM} R \iff \operatorname{Hom}_R(M,X) \in \operatorname{CM} R$$

and

$$\operatorname{Hom}_{\Lambda_M}(\operatorname{Hom}_R(M,X), \Lambda_M) \in \operatorname{CM} R \iff \operatorname{Hom}_{\Lambda_M}(M,X) \in \operatorname{CM} R.$$

Thus there exists $X \in \operatorname{ref} R$ such that $\operatorname{End}_R(M \oplus X) \in \operatorname{CM} R$ if and only if there exists $X \in \operatorname{ref} R$ with $\operatorname{Hom}_R(M,X) \in \operatorname{CM} R$ and $\operatorname{End}_{\Lambda_M}(\operatorname{Hom}_R(M,X)) \in \operatorname{CM} R$. That is, if and only if there exists $Y \in \operatorname{CM}\Lambda_M$ with $\operatorname{End}_{\Lambda_M}(Y) \in \operatorname{CM}(\Lambda_M)$. The statement now follows from the depth lemma (see e.g. [IW10, 2.5]).
(2) Is immediate from (1), since if Λ_M has isolated singularities then $\operatorname{Ext}^1_{\Lambda_M}(Y,Y)$ has finite length. $\square$

Remark 2.18. We conjectured in [IW10] that in dimension three MMAs always have isolated singularities (see 3.13 for some evidence), so the key property from 2.17 is that the stable category of CM modules is rigid-free. Note that $\operatorname{End}_R(M)$ is a NCCR if and only if the stable category is zero. Hence when passing from NCCRs to MMAs, the categories under consideration are no longer zero, but instead are replaced with categories that are rigid-free. This motivates §3.

One application of the results so far is the following much simpler proof of [IW10, 4.14], which avoids many subtle global–local arguments. We will use the following later.

Proposition 2.19. *Suppose that R is normal 3-CY$^-$, and $M \in \operatorname{ref} R$ is a modifying R-module. Suppose that $N \in \operatorname{ref} R$ such that $\operatorname{End}_R(N)$ is derived equivalent to $\operatorname{End}_R(M)$. Then*
(1) N is also modifying.
(2) If M is maximal modifying, so is N.
(3) If M is CT, so is N.

Proof. (1) Since $\operatorname{End}_R(M)$ is 3-CY$^-$ by 2.10, so is $\operatorname{End}_R(N)$ since 3-CY$^-$ algebras are closed under derived equivalences. Hence $\operatorname{End}_R(N) \in \operatorname{CM} R$ by 2.10.
(2) The derived equivalence induces an equivalence between the singular derived categories by 2.7. Since both M and N are modifying by (1), by 2.3 we obtain

$$\underline{\operatorname{CM}} \operatorname{End}_R(M) \approx \operatorname{D_{sg}}(\operatorname{End}_R(M)) \approx \operatorname{D_{sg}}(\operatorname{End}_R(N)) \approx \underline{\operatorname{CM}} \operatorname{End}_R(N).$$

Hence the result follows from 2.17(1).
(3) Since finite global dimension is preserved under derived equivalences, this is obvious [IW10, 2.19]. □

3. Singular Derived Categories and $\mathbb{Q}$-factorial terminalizations

3.1. **Preliminaries.** Recall that a scheme X is called *Gorenstein* if $\mathcal{O}_{X,x}$ is a Gorenstein local ring for all closed points $x \in X$. We say that a scheme X satisfies (ELF) if X is separated, noetherian, of finite Krull dimension, such that $\operatorname{coh} X$ has enough locally free sheaves. This is automatic in many situations, for example if X is quasi–projective.

Definition 3.1. *Suppose that X is a Gorenstein scheme. We say that $\mathcal{F} \in \operatorname{coh} X$ is a* Cohen–Macaulay *(=CM) sheaf if $\mathcal{E}xt^i(\mathcal{F}, \mathcal{O}_X) = 0$ for all $i > 0$.*

Under the assumption that X is Gorenstein, $\mathcal{F} \in \operatorname{coh} X$ is a CM sheaf if and only if

$$0 = \mathcal{E}xt^i(\mathcal{F}, \mathcal{O}_X)_x = \operatorname{Ext}^i_{\mathcal{O}_{X,x}}(\mathcal{F}_x, \mathcal{O}_{X,x})$$

for all $x \in X$ and all $i > 0$, i.e. if and only if $\mathcal{F}_x$ is a CM $\mathcal{O}_{X,x}$-module for all $x \in X$.

Lemma 3.2. *Let R be a noetherian ring of finite Krull dimension, then for any $M \in \operatorname{Mod} R$*

$$\operatorname{proj.dim}_{R_\mathfrak{m}}(M_\mathfrak{m}) < \infty \text{ for all } \mathfrak{m} \in \operatorname{Max} R \iff \operatorname{proj.dim}_R(M) \leq \dim R.$$

In particular, if R has isolated singularities $\{\mathfrak{m}_1, \ldots, \mathfrak{m}_n\}$, then for any $M \in \operatorname{Mod} R$

$$\operatorname{proj.dim}_{R_{\mathfrak{m}_i}}(M_{\mathfrak{m}_i}) < \infty \text{ for all } 1 \leq i \leq n \iff \operatorname{proj.dim}_R(M) \leq \dim R.$$

Proof. We prove the first statement, since the second follows from the first.
($\Leftarrow$) is trivial
($\Rightarrow$) By a result of Bass and Gruson–Raynaud [GR71, 3.2.6] (see also [F77, 3.2]), for any commutative noetherian ring A,

$$\operatorname{FPD}(A) = \sup\{\operatorname{proj.dim}(M) : M \in \operatorname{Mod} A \text{ and } \operatorname{proj.dim}_A(M) < \infty\} = \dim A.$$

The hypothesis implies that $\operatorname{proj.dim}_{R_\mathfrak{p}}(M_\mathfrak{p}) < \infty$ for all $\mathfrak{p} \in \operatorname{Spec} R$, thus by the above, necessarily $\operatorname{proj.dim}_{R_\mathfrak{p}}(M_\mathfrak{p}) \leq \dim R_\mathfrak{p} \leq \dim R$ for all $\mathfrak{p} \in \operatorname{Spec} R$. In particular $\operatorname{flat.dim}_{R_\mathfrak{p}}(M_\mathfrak{p}) \leq \dim R$ for all $\mathfrak{p} \in \operatorname{Spec} R$. Now $X \in \operatorname{Mod} R$ is zero if $X_\mathfrak{p} = 0$ for all $\mathfrak{p} \in \operatorname{Spec} R$, so since Tor groups always localize, M has finite flat dimension. Thus by [J70, Prop 6] M has finite projective dimension, which by the result of Bass and Gruson–Raynaud above is necessarily less than or equal to $\dim R$. □

Definition 3.3. *We say that $\mathcal{F} \in \operatorname{Qcoh} X$ is locally free if X can be covered with open sets U for which each $\mathcal{F}|_U$ is a (not necessarily finitely generated) free $\mathcal{O}_X|_U$-module.*

As in [Or03, §1] we consider

- the full triangulated subcategory $\mathfrak{Perf}(X)$ of $\mathrm{D}^{\mathrm{b}}(\operatorname{coh} X)$ consisting of objects which are isomorphic to bounded complexes of finitely generated locally free sheaves in $\mathrm{D}^{\mathrm{b}}(\operatorname{coh}(X))$.
- the full triangulated subcategory $\mathfrak{Lfr}(X)$ of $\mathrm{D}^{\mathrm{b}}(\operatorname{Qcoh} X)$ consisting of objects which are isomorphic to bounded complexes of locally free sheaves in $\mathrm{D}^{\mathrm{b}}(\operatorname{Qcoh}(X))$.

Following Orlov [Or03, Or09], we denote $\mathrm{D}_{\mathsf{sg}}(X) := \mathrm{D}^{\mathrm{b}}(\operatorname{coh} X)/\,\mathfrak{Perf}(X)$ and $\mathrm{D}'_{\mathsf{sg}}(X) := \mathrm{D}^{\mathrm{b}}(\operatorname{Qcoh} X)/\,\mathfrak{Lfr}(X)$. The following is evident, and is well-known:

Lemma 3.4. *Let R be a commutative noetherian ring of finite Krull dimension, then* $\mathrm{K}^{\mathrm{b}}(\operatorname{Free} R) \approx \mathfrak{Lfr}(\operatorname{Spec} R) \approx \mathrm{K}^{\mathrm{b}}(\operatorname{Proj} R)$.

Proof. We have natural inclusions $\mathrm{K}^{\mathrm{b}}(\operatorname{Free} R) \subseteq \mathfrak{Lfr}(\operatorname{Spec} R) \subseteq \mathrm{K}^{\mathrm{b}}(\operatorname{Proj} R)$ since every free module is clearly locally free, and further every locally free R-module is locally projective, thus projective by a result of Gruson–Raynaud [GR71, 3.1.4(3)]. We claim that $\mathrm{K}^{\mathrm{b}}(\operatorname{Proj} R) = \mathrm{K}^{\mathrm{b}}(\operatorname{Free} R)$, as then we have equality throughout. This is similar to [R89, 2.2] — by the Eilenberg swindle, if $P \in \operatorname{Proj} R$, there exists $F \in \operatorname{Free} R$ such that $P \oplus F$ is free. Hence from every object of $\mathrm{K}^{\mathrm{b}}(\operatorname{Proj} R)$ we can get to an object of $\mathrm{K}^{\mathrm{b}}(\operatorname{Free} R)$ by taking the direct sum with complexes of the form $0 \to F \to F \to 0$, which are zero objects in the homotopy category. ☐

Corollary 3.5. *Suppose X satisfies (ELF), and has isolated singularities $\{x_1, \dots, x_n\}$. If $\mathcal{F} \in \operatorname{Qcoh} X$ with $\operatorname{proj.dim}_{\mathcal{O}_{X,x_i}}(\mathcal{F}_{x_i}) < \infty$ for all $1 \leq i \leq n$, then $\mathcal{F} \in \mathfrak{Lfr}(X)$.*

Proof. Choose an affine cover of X, then on each affine piece $U = \operatorname{Spec} R$, $\mathcal{F}_{\mathfrak{m}}$ is an $R_{\mathfrak{m}}$-module with finite projective dimension for all the isolated singularities $\mathfrak{m}$ lying in U. Hence $\mathcal{F}|_U \in \mathrm{K}^{\mathrm{b}}(\operatorname{Proj} R)$ by 3.2, so by 3.4 it follows that $\mathcal{F}|_U$ is isomorphic, in $\mathrm{D}^{\mathrm{b}}(\operatorname{Qcoh} U)$, to a complex in $\mathfrak{Lfr}(U)$. Since X satsifies (ELF), membership of $\mathfrak{Lfr}(X)$ can be checked locally (see [Or03, 1.7, proof of 1.14]), so it follows that $\mathcal{F} \in \mathfrak{Lfr}(X)$. ☐

We will also require the following well-known lemma, due to Orlov. If $\mathcal{S}$ is an object in a triangulated category $\mathcal{T}$, we denote $\langle \mathcal{S} \rangle_{\mathcal{T}}$ to be the smallest thick triangulated subcategory of $\mathcal{T}$ containing $\mathcal{S}$.

Lemma 3.6. (Orlov) *Suppose that X satisfies (ELF), and has only isolated singularities $\{x_1, \dots, x_n\}$. Denote the corresponding skyscraper sheaves by $k_1, \dots, k_n$. Then $\mathrm{D}_{\mathsf{sg}}(X) = \langle \oplus_{i=1}^n k_i \rangle_{\mathrm{D}_{\mathsf{sg}}(X)}$.*

Proof. By assumption the singular locus consists of a finite number of closed points, hence is closed. It follows that $\mathrm{D}_{\mathsf{sg}}(X) = \langle \operatorname{coh}_{\{x_1,\dots,x_n\}} X \rangle_{\mathrm{D}_{\mathsf{sg}}(X)}$ by [C10, 1.2]. It is clear that every coherent sheaf supported in $\{x_1, \dots, x_n\}$ belongs to $\langle \oplus_{i=1}^n k_i \rangle_{\mathrm{D}_{\mathsf{sg}}(X)}$. ☐

The following is “folklore” in the sense that many people expected it to be true, however in the non-local case subtleties arise in the proof. Certainly the idempotent completion part also follows from work of Orlov [Or09], and is well-known to experts [BK11]:

Theorem 3.7. *Suppose that X is a Gorenstein scheme of dimension d satisfying (ELF), with isolated singularities $\{x_1, \dots, x_n\}$.*

(1) *There is a fully faithful functor* $\mathrm{D}_{\mathsf{sg}}(X) \hookrightarrow \displaystyle\bigoplus_{i=1}^{n} \underline{\mathrm{CM}}\, \mathcal{O}_{X,x_i}$ *which is essentially surjective up to summands.*

(2) *Taking the idempotent completion gives an equivalence* $\overline{\mathrm{D}_{\mathsf{sg}}(X)} \approx \bigoplus_{i=1}^{n} \underline{\mathrm{CM}}\, \widehat{\mathcal{O}}_{X,x_i}$.

Proof. For each x_i, consider a morphism $f_i := \operatorname{Spec} \mathcal{O}_{X,x_i} \xrightarrow{g_i} \operatorname{Spec} R_i \xrightarrow{h_i} X$ where $\operatorname{Spec} R_i$ is some affine open containing x_i. The functor $g_{i\,*}$ is just extension of scalars corresponding to a localization $R_i \to R_{i\mathfrak{m}}$, so $g_{i\,*}$ is exact and preserves quasi-coherence. Further h_i is an affine morphism, hence $h_{i\,*}$ also is exact and preserves quasi-coherence. Hence each $f_{i\,*}$ is exact and preserves quasi-coherence.

Now the collection of the f_i induce a morphism

$$\operatorname{Spec}(\mathcal{O}_{X,x_1} \oplus \ldots \oplus \mathcal{O}_{X,x_n}) = \coprod \operatorname{Spec} \mathcal{O}_{X,x_i} \xrightarrow{f} X.$$

For convenience, denote $S := \mathcal{O}_{X,x_1} \oplus \ldots \oplus \mathcal{O}_{X,x_n}$, then the above induces an adjunction

$$\begin{tikzcd} \operatorname{Qcoh} X \arrow[r, shift left, "f^*"] & \operatorname{Qcoh} S = \bigoplus_{i=1}^{n} \operatorname{Qcoh} \mathcal{O}_{X,x_i} \arrow[l, shift left, "f_*"] \end{tikzcd}$$

which explicitly is given by $f^*\mathcal{F} = (\mathcal{F}_{x_1}, \ldots, \mathcal{F}_{x_n})$ and $f_*(\mathcal{G}_1, \ldots, \mathcal{G}_n) = \oplus f_{i*}\mathcal{G}_i$. These functors are exact, so induce an adjunction

$$\begin{tikzcd} \mathrm{D^b}(\operatorname{Qcoh} X) \arrow[r, shift left, "f^*"] & \mathrm{D^b}(\operatorname{Qcoh} S) = \bigoplus_{i=1}^{n} \mathrm{D^b}(\operatorname{Qcoh} \mathcal{O}_{X,x_i}) \arrow[l, shift left, "f_*"] \end{tikzcd}$$

in the obvious way. It is clear that f^* takes locally free sheaves to projective modules, and so sends $\mathfrak{Lfr}(X)$ to $\mathrm{K^b}(\operatorname{Proj} S) = \mathfrak{Lfr}(S)$.

Now each f_{i*} is an affine, flat morphism so $f_{i*}\mathcal{O}_{X,x_i}$ is a flat $\mathcal{O}_X$-module. Further, each f_{i*} preserves sums, so takes projective $\mathcal{O}_{X,x_i}$-modules to flat $\mathcal{O}_X$-modules. Consequently, it follows that f_* takes a projective S-module P to a sheaf $f_*(P)$ for which $f_*(P)_x$ is a flat $\mathcal{O}_{X,x}$-module for all $x \in X$. Since flat modules have finite projective dimension (by [J70, Prop 6]) certainly $\operatorname{proj.dim}_{\mathcal{O}_{X,x_i}}(f_*(P)_{x_i}) < \infty$ for all $1 \leq i \leq n$, hence by 3.5 $f_*(P) \in \mathfrak{Lfr}(X)$. This implies that f_* takes a complex C in $\mathrm{K^b}(\operatorname{Proj} S) = \mathfrak{Lfr}(S)$ to a bounded complex $f_*(C)$ which has, in each non-zero degree, an object in $\mathfrak{Lfr}(X)$. Since $\mathfrak{Lfr}(X)$ is a full triangulated subcategory, by induction on the number of non-zero terms it follows that $f_*(C)$ belongs to $\mathfrak{Lfr}(X)$.

Hence the above exact adjunction induces an exact adjunction

$$\begin{tikzcd} \mathrm{D}'_{\mathsf{sg}}(X) \arrow[r, shift left, "f^*"] & \mathrm{D}'_{\mathsf{sg}}(S) = \bigoplus_{i=1}^{n} \mathrm{D}'_{\mathsf{sg}}(\mathcal{O}_{X,x_i}). \arrow[l, shift left, "f_*"] \end{tikzcd}$$

Now denote the skyscraper sheaves in X corresponding to the singular points by $k_1, \ldots, k_n$. By 3.6 we know that the thick closure (taken inside $\mathrm{D}_{\mathsf{sg}}(X)$) of $\oplus_{i=1}^{n} k_i$ equals $\mathrm{D}_{\mathsf{sg}}(X)$. We do not necessarily know that $\mathrm{D}_{\mathsf{sg}}(X)$ is thick inside $\mathrm{D}'_{\mathsf{sg}}(X)$, however certainly $\langle \oplus_{i=1}^{n} k_i \rangle_{\mathrm{D}'_{\mathsf{sg}}}$, the thick closure of $\oplus_{i=1}^{n} k_i$ inside $\mathrm{D}'_{\mathsf{sg}}(X)$, contains $\mathrm{D}_{\mathsf{sg}}(X)$.

Note that $f_* \circ f^*(k_i) = f_*(0, \ldots, 0, k_i, 0, \ldots, 0) = f_{i*}k_i = k_i$ for all $1 \leq i \leq n$. This implies that $f_* \circ f^* \cong 1$ on $\langle \oplus_{i=1}^{n} k_i \rangle_{\mathrm{D}'_{\mathsf{sg}}}$, and so f^* is fully faithful out of $\langle \oplus_{i=1}^{n} k_i \rangle_{\mathrm{D}'_{\mathsf{sg}}}$. It is clear that f^* takes $\mathrm{D}_{\mathsf{sg}}(X)$ to $\mathrm{D}_{\mathsf{sg}}(S)$, so since $\mathrm{D}_{\mathsf{sg}}(X)$ is contained inside $\langle \oplus_{i=1}^{n} k_i \rangle_{\mathrm{D}'_{\mathsf{sg}}}$, we deduce that

$$\mathrm{D}_{\mathsf{sg}}(X) \xrightarrow{f^*} \mathrm{D}_{\mathsf{sg}}(S)$$

is fully faithful. Now $\operatorname{Im} f^*$ is a triangulated subcategory of $\mathrm{D}_{\mathsf{sg}}(S)$ and by the calculation above $\operatorname{Im} f^*$ contains the set $\mathcal{S} := \{(0, \ldots, 0, k_i, 0, \ldots, 0) \mid 1 \leq i \leq n\}$. By 3.6 we know that these generate the components $\mathrm{D}_{\mathsf{sg}}(\mathcal{O}_{X,x_i})$ of $\mathrm{D}_{\mathsf{sg}}(S)$, That is to say, the thick closure (inside $\mathrm{D}_{\mathsf{sg}}(S)$) of the set $\mathcal{S}$ equals $\mathrm{D}_{\mathsf{sg}}(S)$. Now $\operatorname{Im} f^*$ is a triangulated category containing $\mathcal{S}$, hence the thick closure of $\operatorname{Im} f^*$ equals $\mathrm{D}_{\mathsf{sg}}(S)$. But always, the thick closure of a triangulated subcategory is the class of summands of objects of that subcategory [N01, 2.1.39], so we deduce that f^* is an equivalence up to summands.

The first statement in the theorem now follows from the well-known equivalence $\mathrm{D}_{\mathsf{sg}}(\mathcal{O}_{X,x_i}) \approx \underline{\mathrm{CM}}\,\mathcal{O}_{X,x_i}$ due to Buchweitz [Bu86, 4.4.1]. The second statement follows from (1), since the idempotent completion of $\underline{\mathrm{CM}}\,\mathcal{O}_{X,x_i}$ is $\underline{\mathrm{CM}}\,\widehat{\mathcal{O}}_{X,x_i}$ [KMV, A.1]. $\square$

The first corollary of 3.7 is the following alternative proof of [Or03, 1.24], which we will use later:

Corollary 3.8. *Suppose that X is a d-dimensional Gorenstein scheme over k, satisfying (ELF), with isolated singularities. Then all hom-spaces in $\mathrm{D}_{\mathsf{sg}}(X)$ have finite k-dimension.*

Proof. Since each $\mathcal{O}_{X,x_i}$ is isolated, $\operatorname{Hom}_{\underline{\mathrm{CM}}\,\mathcal{O}_{X,x_i}}(A, B) = \operatorname{Ext}^1_{\mathcal{O}_{X,x_i}}(\Omega A, B)$ is supported only on the maximal ideal of $\mathcal{O}_{X,x_i}$, hence has finite length. But each $\mathcal{O}_{X,x_i}$ is a finitely generated k-algebra, so by the Nullstellensatz its residue field is a finite extension of k. By 3.7(1), the result follows. $\square$

3.2. **Rigid-free, $\mathbb{Q}$-factoriality and MMAs.** In this section we characterize the $\mathbb{Q}$-factorial property in terms of the singular derived category, and then relate this to MMAs. The following is a generalization of 2.13(3)⇒(4).

Proposition 3.9. *Suppose that X is a Gorenstein normal scheme satisfying (ELF), of dimension three which has only isolated singularities $\{x_1, \dots, x_n\}$. If $\mathrm{D_{sg}}(X)$ is rigid-free, then X is locally factorial.*

Proof. Suppose that $\mathrm{D_{sg}}(X)$ is rigid-free. By Auslander–Buchsbaum, any CM sheaf which is zero in $\mathrm{D_{sg}}(X)$ is necessarily locally free. Hence the assumption implies that if $\mathcal{F}$ is some CM sheaf on X for which $\mathrm{Hom}_{\mathrm{D_{sg}}(X)}(\mathcal{F}, \mathcal{F}[1]) = 0$, then $\mathcal{F}$ is locally free. But by 3.7(1)

$$\mathrm{Hom}_{\mathrm{D_{sg}}(X)}(\mathcal{F}, \mathcal{F}[1]) \cong \bigoplus_{i=1}^{n} \mathrm{Hom}_{\mathrm{D_{sg}}(\mathcal{O}_{X,x_i})}(\mathcal{F}_{x_i}, \mathcal{F}_{x_i}[1]) \cong \bigoplus_{i=1}^{n} \mathrm{Ext}^1_{\mathcal{O}_{X,x_i}}(\mathcal{F}_{x_i}, \mathcal{F}_{x_i}),$$

hence if $\mathcal{F}$ is a CM sheaf on X for which $\mathrm{Ext}^1_{\mathcal{O}_{X,x_i}}(\mathcal{F}_{x_i}, \mathcal{F}_{x_i}) = 0$ for all $1 \le i \le n$, then $\mathcal{F}$ is locally free.

Now let $\mathcal{G} \in \mathrm{Cl}(X)$. Certainly $\mathcal{G}_x \in \mathrm{Cl}(\mathcal{O}_{X,x})$ for all $x \in X$, so since $\mathcal{O}_{X,x}$ is normal we have $\mathrm{End}_{\mathcal{O}_{X,x}}(\mathcal{G}_x) \cong \mathcal{O}_{X,x} \in \mathrm{CM}\,\mathcal{O}_{X,x}$ for all $x \in X$. But $\mathcal{O}_{X,x}$ is isolated, so by [IW10, 2.6] $\mathrm{Ext}^1_{\mathcal{O}_{X,x}}(\mathcal{G}_x, \mathcal{G}_x) = 0$ for all $x \in X$. Since X satisfies (ELF) we may find a short exact sequence

$$0 \to \mathcal{K} \to \mathcal{V} \to \mathcal{G} \to 0 \tag{3.D}$$

with $\mathcal{V}$ a locally free sheaf. Now localizing (3.D) to a point $x \in X$ gives

$$0 \to \mathcal{K}_x \to \mathcal{V}_x \to \mathcal{G}_x \to 0 \tag{3.E}$$

and so since $\mathcal{G}_x \in \mathrm{ref}\,\mathcal{O}_{X,x}$, by counting depths we see that $\mathcal{K}_x \in \mathrm{CM}\,\mathcal{O}_{X,x}$. This holds for all $x \in X$, so $\mathcal{K}$ is a CM sheaf. But on the other hand (3.E) is an $\mathcal{O}_{X,x}$-approximation of the reflexive $\mathcal{O}_{X,x}$-module $\mathcal{G}_x$, so by [IW10, 4.4] $\mathrm{End}_{\mathcal{O}_{X,x}}(\mathcal{K}_x) \in \mathrm{CM}\,\mathcal{O}_{X,x}$. Since $\mathcal{O}_{X,x}$ is isolated, this forces $\mathrm{Ext}^1_{\mathcal{O}_{X,x}}(\mathcal{K}_x, \mathcal{K}_x) = 0$ for all $x \in X$. Since $\mathcal{K}$ is a CM sheaf, by above we deduce that $\mathcal{K}$ must be locally free.

Thus (3.E) now shows that $\mathrm{proj.dim}_{\mathcal{O}_{X,x}} \mathcal{G}_x \le 1$ for all $x \in X$. By [AG60, 4.10], if $\mathrm{proj.dim}_{\mathcal{O}_{X,x}} \mathcal{G}_x = 1$ then $\mathrm{Ext}^1_{\mathcal{O}_{X,x}}(\mathcal{G}_x, \mathcal{G}_x) \neq 0$. Since $\mathrm{Ext}^1_{\mathcal{O}_{X,x}}(\mathcal{G}_x, \mathcal{G}_x) = 0$, it follows that $\mathrm{proj.dim}_{\mathcal{O}_{X,x}} \mathcal{G}_x = 0$. This shows that $\mathcal{G}$ is locally free, hence every member of the class group is locally free and so X is locally factorial. ☐

Even although we do not know if the functor in 3.7(1) is strictly an equivalence, provided the singularities are at worst hypersurfaces, the existence of rigid objects can be checked on either side:

Theorem 3.10. *Suppose that X is a normal 3-dimensional Gorenstein scheme over k, satisfying (ELF), with isolated singularities $\{x_1, \dots, x_n\}$.*

(1) *If $\mathcal{O}_{X,x_i}$ are hypersurfaces for all $1 \le i \le n$, then the following are equivalent:*
   (a) *X is locally factorial.*
   (b) *X is $\mathbb{Q}$-factorial.*
   (c) *$\mathrm{D_{sg}}(X)$ is rigid-free.*
   (d) *$\underline{\mathrm{CM}}\,\mathcal{O}_{X,x}$ is rigid-free for all closed points $x \in X$.*

(2) *If $\widehat{\mathcal{O}}_{X,x_i}$ are hypersurfaces for all $1 \le i \le n$, then the following are equivalent:*
   (a) *X is complete locally factorial.*
   (b) *X is complete locally $\mathbb{Q}$-factorial.*
   (c) *$\overline{\mathrm{D_{sg}}(X)}$ is rigid-free.*
   (d) *$\underline{\mathrm{CM}}\,\widehat{\mathcal{O}}_{X,x}$ is rigid-free for all closed points $x \in X$.*

Proof. (1) (a)⇒(b) is clear.
(b)⇒(d) follows from 2.14.
(d)⇒(c) follows from 3.7(a).
(c)⇒(a) follows from 3.9.
(2) The proof is identical to the proof in (1). ☐

We define ‘derived equivalence’ on the bounded level, so we say that X is *derived equivalent* to Λ if and only if $\mathrm{D^b}(\mathrm{coh}\, X)$ is equivalent to $\mathrm{D^b}(\mathrm{mod}\, \Lambda)$.

Lemma 3.11. *Suppose that X satisfies (ELF), and is over a field k. Suppose that Λ is a module-finite R-algebra, such that X is derived equivalent to Λ. Then*
(1) $\mathfrak{Perf}(X) \approx \mathrm{K^b}(\mathrm{proj}\, \Lambda)$.
(2) $\mathrm{D_{sg}}(X) \approx \mathrm{D_{sg}}(\Lambda)$.
(3) *X is smooth* $\iff$ $\mathrm{gl.dim}\, \Lambda < \infty$.

Proof. (1) In [Or05, 1.11], under the assumptions above Orlov shows that $\mathfrak{Perf}(X)$ are precisely the homologically finite complexes in $\mathrm{D^b}(\mathrm{coh}\, X)$. The result follows by 2.7.
(2) follows from (1).
(3) We first claim that X is smooth $\iff$ $\mathrm{D_{sg}}(X) = 0$. This is well known — certainly if X is smooth then $\mathrm{D_{sg}}(X) = 0$ [Or03, 1.9]. If $\mathrm{D_{sg}}(X) = 0$, then every coherent sheaf has finite homological dimension, hence the skyscrapers all have finite homological dimension, hence X is smooth. Since Λ is noetherian, it is also well know that $\mathrm{D_{sg}}(\Lambda) = 0$ if and only if $\mathrm{gl.dim}\, \Lambda < \infty$ [R08, 7.26]. Hence the result follows by (2). ☐

Corollary 3.12. *Let R be normal d-CY$^-$ over k, and suppose that $M \in \mathrm{ref}\, R$ is modifying. Suppose that Λ is derived equivalent to X, where X is some d-dimensional Gorenstein scheme, over k, satisfying (ELF), with only isolated singularities. Then Λ has isolated singularities.*

Proof. The derived equivalence induces $\mathrm{D_{sg}}(X) \approx \mathrm{D_{sg}}(\Lambda)$ by 3.11, hence the result follows by combining 3.8 and 2.4. ☐

We are now in a position to relate maximal modification algebras and $\mathbb{Q}$-factorial terminalizations. The following is the main result of this subsection:

Theorem 3.13. *Let R be 3-dimensional normal Gorenstein domain over k, and suppose that $Y \to \mathrm{Spec}\, R$ is a proper birational map, where Y is normal and has only isolated singularities $\{x_1, \dots, x_n\}$ where each $\mathcal{O}_{Y,x_i}$ is a hypersurface. Assume that Y is derived equivalent to $\mathrm{End}_R(M)$ for some modifying R-module M, then*
(1) Y is $\mathbb{Q}$-factorial $\iff$ $\mathrm{End}_R(M)$ is a MMA with isolated singularities.
(2) All maximal modification algebras have isolated singularities, and are all derived equivalent.

Proof. Since the singularities are isolated, note first that $\Lambda := \mathrm{End}_R(M)$ has isolated singularities by 3.12.
(1) The derived equivalence forces $\mathrm{D_{sg}}(Y) \approx \mathrm{D_{sg}}(\Lambda)$ by 3.11, which by 2.3 is equivalent to $\underline{\mathrm{CM}}\, \Lambda$. The result follows by combining 2.17 and 3.10(1).
(2) Since Λ has isolated singularities, all maximal modification algebras have isolated singularities and are derived equivalent by [IW10, 4.9, 4.12]. ☐

Remark 3.14. When $k = \mathbb{C}$ and Y has only terminal singularities, the geometric assumptions in 3.13 are satisfied. Terminal singularities are isolated for 3-folds [KM98, 5.18], and Gorenstein terminal singularities are Zariski locally hypersurfaces [R83, 0.6(I)]

The following corollary will be used in §4.

Corollary 3.15. *With the setup as in 3.13, suppose further that R is complete local. Then the following are equivalent.*
(1) *X is $\mathbb{Q}$-factorial.*
(2) *$\mathrm{End}_R(M)$ is a MMA with isolated singularities.*
(3) *X is complete locally $\mathbb{Q}$-factorial.*

Proof. (1)⇒(2) is 3.13(1).
(2)⇒(3) Since R is complete, $\underline{\mathrm{CM}}\, \mathrm{End}_R(M)$ is already idempotent complete, since it is an additive category which is Krull–Schmidt [CYZ, A.1]. This implies that $\mathrm{D_{sg}}(\mathrm{End}_R(M)) = \overline{\mathrm{D_{sg}}(\mathrm{End}_R(M))}$ and so the result follows by 3.10(2).
(3)⇒(1) is obvious. ☐

4. cA_n SINGULARITIES VIA DERIVED CATEGORIES

We now illustrate the results of the previous section to give many examples of maximal modification algebras.

Proposition 4.1. (Shepherd–Barron, unpublished) *Let* $R = \mathbb{C}[u, v, x, y]/uv - f(x, y)$ *be an isolated* cA_n *singularity. Then the following are equivalent*
(1) *There does not exists a non-trivial crepant morphism* $Y \to \operatorname{Spec} R$.
(2) $f(x, y)$ *is irreducible.*
(3) R *is factorial.*
(4) R *is a maximal modifying* R*-module.*

Proof. (1)⇒(2) If $f(x, y)$ factors as $f = f_1 f_2$ then blowing up the ideal (u, f_1) yields a non-trivial crepant morphism. Hence $f(x, y)$ cannot factor.
(2)⇒(3) is clear (see e.g. proof of 5.4(1)).
(3)⇒(1) If there exists such a non-trivial crepant map, since R is terminal necessarily the map must have one-dimensional fibres. Hence by [V04a, 3.2.10] there exists a non-trivial modifying module, so it follows from 2.14 that R cannot be factorial, a contradiction.
(3)⇔(4) is a special case of 2.15. ☐

4.1. **Crepant Modifications of cA_n singularities.** In this section we work over the field $\mathbb{K}$ and restrict to the complete local case, since it vastly simplifies the proof of the main theorem 4.2. See 4.7 for remarks on the non-local case.

Define $R := \mathbb{K}[[u, v, x, y]]/(uv - f(x, y))$, with $f \in \mathfrak{m} := (x, y) \subseteq \mathbb{K}[[x, y]]$. We let $f = f_1 \dots f_n$ be a factorization into (not necessarily distinct) prime elements of $\mathbb{K}[[x, y]]$. We remark that when $\mathbb{K} = \mathbb{C}$, R is a cDV singularity of type A (see e.g. [BIKR, 6.1(e)]), which is terminal if and only if R is an isolated singularity [R83, 1.1].

For any subset $I \subseteq \{1, \dots, n\}$ we denote

$$f_I := \prod_{i \in I} f_i \quad \text{and} \quad T_I := (u, f_I)$$

which is an ideal of R. For a collection of subsets

$$\emptyset \subsetneq I_1 \subsetneq I_2 \subsetneq \dots \subsetneq I_m \subsetneq \{1, 2, \dots, n\},$$

we say that $\mathcal{F} = (I_1, \dots, I_m)$ is a *flag in the set* $\{1, 2, \dots, n\}$. We say that the flag $\mathcal{F}$ is *maximal* if $n = m + 1$. Given a flag $\mathcal{F} = (I_1, \dots, I_m)$, we define

$$T^{\mathcal{F}} := R \oplus \left(\bigoplus_{j=1}^{m} T_{I_j} \right).$$

Geometrically, given a flag $\mathcal{F} = (I_1, \dots, I_m)$ we define a space $X^{\mathcal{F}}$ as follows: first, blowup the ideal (u, f_{I_1}) in $R = \mathbb{K}[[u, v, x, y]]/(uv - f)$. This yields a space X^{I_1} covered by two open charts, given explicitly by

$$R_1 := \frac{\mathbb{K}[[u, x, y]][V]}{uV = f_{I_1}} \quad \text{and} \quad R_2 := \frac{\mathbb{K}[[v, x, y]][U]}{Uv = \frac{f}{f_{I_1}}}.$$

Note that the new co-ordinates $V = \frac{f_{I_1}}{u}$ and $U = \frac{u}{f_{I_1}}$ glue to give a copy of $\mathbb{P}^1$ inside X^{I_1} (possibly moving in a family), which maps to the origin of $\operatorname{Spec} R$.

Next blowup the divisor (U, f_{I_2}) in the second co-ordinate chart R_2, and call this space X^{I_2}. Note that the zero set of the divisor (U, f_{I_2}) does not intersect the first co-ordinate chart R_1, so under the second blowup R_1 is unaffected. Locally above R_2, the calculation to determine the structure of X^{I_2} is identical to the above, giving a cover of X^{I_2} by the three affine open sets

$$\frac{\mathbb{K}[[u, x, y]][V]}{uV = f_{I_1}} \quad \text{and} \quad \frac{\mathbb{K}[[u, x, y]][V]}{uV = f_{I_2}} \quad \text{and} \quad \frac{\mathbb{K}[[v, x, y]][U]}{Uv = \frac{f}{f_{I_1} f_{I_2}}}.$$

Hence we have produced another $\mathbb{P}^1$ (again which might move in a family) which maps to the origin of R_2 and hence to the origin of R.

Continuing by blowing up the ideal (U, f_{I_3}), in this way we obtain a chain of projective birational morphisms

$$X^{I_m} \to X^{I_{m-1}} \to \ldots \to X^{I_1} \to \operatorname{Spec} R,$$

and we define $X^{\mathcal{F}} := X^{I_m}$. See 4.6 later for a picture of this process.

Theorem 4.2. *Given a flag $\mathcal{F} = (I_1, \ldots, I_m)$, denote $X^{\mathcal{F}}$ and $T^{\mathcal{F}}$ as above. Then $X^{\mathcal{F}}$ is derived equivalent to $\operatorname{End}_R(T^{\mathcal{F}})$.*

Proof. In the explicit calculation for $X^{\mathcal{F}}$ above, the preimage of the unique closed point of $\operatorname{Spec} R$ is a chain of $\mathbb{P}^1$'s (some of which move in a family), in a type-A configuration. Now by [V04a, Thm. B], there is a tilting bundle on $X^{\mathcal{F}}$ given as follows: let $C = f^{-1}(\mathfrak{m})$ where $\mathfrak{m}$ is the unique closed point of $\operatorname{Spec} R$. Giving C the reduced scheme structure, write $C_{\text{red}} = \cup_{i \in I} C_i$, and let $\mathcal{L}_i$ denote the line bundle on $X^{\mathcal{F}}$ such that $\mathcal{L}_i \cdot C_j = \delta_{ij}$. If the multiplicity of C_i in C is equal to one, set $\mathcal{M}_i := \mathcal{L}_i$ [V04a, 3.5.4], else define $\mathcal{M}_i$ to be given by the extension

$$0 \to \mathcal{O}^{r_i - 1} \to \mathcal{M}_i \to \mathcal{L}_i \to 0$$

associated to a minimal set $r_i - 1$ of generators of $H^1(X^{\mathcal{F}}, \mathcal{L}_i^{-1})$. Then $\mathcal{O} \oplus_{i \in I} \mathcal{M}_i$ is a tilting bundle on $X^{\mathcal{F}}$ [V04a, 3.5.5].

In our situation, let C_i be the curve in $X^{\mathcal{F}}$, above the origin of $\operatorname{Spec} R$, which in the process of blowing up first appears in X^{I_i} (see 4.6 for a picture of this). We now claim that all the curves C_i have multiplicity one. If some C_i had multiplicity greater than one, then $\mathcal{M}_i$ would be a indecomposable bundle with rank greater than one [V04a, 3.5.4]. But this would imply, by [V04a, 3.2.9], that its sections $H^0(\mathcal{M}_i) := M_i$ is an indecomposable CM R-module of rank greater than one, such that $\operatorname{End}_R(M_i) \in \operatorname{CM} R$. But this is impossible (see e.g. 5.1). Hence all curves have multiplicity one, and so $\mathcal{O} \oplus_{i=1}^m \mathcal{L}_i$ is a tilting bundle on $X^{\mathcal{F}}$.

Now $\operatorname{End}_{X^{\mathcal{F}}}(\mathcal{O} \oplus_{i=1}^m \mathcal{L}_i) \cong \operatorname{End}_R(R \oplus_{i=1}^m H^0(\mathcal{L}_i))$, and so it remains to show that $R \oplus_{i=1}^m H^0(\mathcal{L}_i) \cong T^{\mathcal{F}}$. In fact, we claim that $H^0(\mathcal{L}_i) \cong (u, f_{I_i})$. But it is easy to see (for example using the Čech complex) that the rank one CM module $H^0(\mathcal{L}_i)$ contains the elements u and f_{I_i}. Hence $H^0(\mathcal{L}_i)$ is a rank one CM module containing the ideal (u, f_{I_i}). But this ideal is a rank one reflexive module (see e.g. 5.4), hence the natural inclusion $(u, f_{I_i}) \to H^0(\mathcal{L}_i)$, being a height one isomorphism, is an isomorphism. □

So as to match our notation with [BIKR] and [DH10], we can (and do) identify maximal flags with elements of the symmetric group $\mathfrak{S}_n$. Hence we regard each $\omega \in \mathfrak{S}_n$ as the maximal flag

$$\{\omega(1)\} \subset \{\omega(1), \omega(2)\} \subset \ldots \subset \{\omega(1), \ldots, \omega(n-1)\}.$$

The following is simply a special case of 4.2.

Corollary 4.3. *Let $\omega \in \mathfrak{S}_n$, then X^ω is derived equivalent to $\operatorname{End}_R(T^\omega)$. Further, X^ω is a space in which the set of completions of the singular points is precisely $\{\mathbb{K}[[u,v,x,y]]/(uv - f_i) \mid 1 \le i \le n, f_i \in (x,y)^2\}$.*

This yields the following generalization of [BIKR, 1.5] and [DH10, 4.2], and provides a conceptual reason for the condition $f_i \notin \mathfrak{m}^2$.

Corollary 4.4. *Suppose $f_1, \ldots, f_n \in \mathfrak{m} := (x,y) \subseteq \mathbb{K}[[x,y]]$ are irreducible polynomials. Let $R = \mathbb{K}[[u,v,x,y]]/(uv - f_1 \ldots f_n)$, then*
(1) Each T^ω ($\omega \in \mathfrak{S}_n$) is a MM R-module generator. The corresponding endomorphism rings all have isolated singularities.
(2) T^ω are CT modules $\iff$ $f_i \notin \mathfrak{m}^2$ for all $1 \le i \le n$.

Proof. (1) $\operatorname{End}_R(T^\omega)$ is derived equivalent to a space whose (complete local) singularities are listed in 4.3. Since each f_i is irreducible, these are all factorial (4.1). Hence the result follows from 3.15.
(2) T^ω is CT if and only if $\operatorname{End}_R(T^\omega)$ is a NCCR [IW10, 5.5]. Since $\operatorname{End}_R(T^\omega) \in \operatorname{CM} R$, this occurs if and only if $\operatorname{gl.dim} \operatorname{End}_R(T^\omega) < \infty$. By 3.11, this occurs if and only if X^ω is smooth. But by 4.3, this happens if and only if each $uv = f_i$ is smooth, which is if and only if each $f_i \notin \mathfrak{m}^2$. □

Remark 4.5. For a non-derived category proof of the above, see 5.17 and 5.19. Note that 5.17 and 5.19 are stronger statements, since they assert that there are no other MM (respectively CT) generators, and they work over more general fields.

Remark 4.6. It is useful to visualize the above in an example. Consider $uv = f_1f_2f_3f_4$ and choose maximal flag $\mathcal{F} = (I_1 := \{1\} \subset I_2 := \{1,2\} \subset I_3 := \{1,2,3\})$. Thus $T_{I_1} = (u, f_1)$, $T_{I_2} = (u, f_1f_2)$ and $T_{I_3} = (u, f_1f_2f_3)$. Then we have the following:


$X^{\mathcal{F}}$
φ_3
X^{I_2}
φ_2
X^{I_1}
φ_1
$\operatorname{Spec} R$
T_{I_1}, T_{I_2}, T_{I_3}, R
inc, f_1, f_2, f_3, $\frac{f_4}{u}$, u, x, y
$\frac{f_3f_4}{u}$
$\frac{f_2f_3f_4}{u}$
v
$R \oplus T_{I_1} \oplus T_{I_2} \oplus T_{I_3}$
$R \oplus T_{I_1} \oplus T_{I_2}$
$R \oplus T_{I_1}$
R

On the geometric side we have drawn the non-local picture; the complete local picture is obtained by drawing a tubular neighbourhood around all the red curves, and restricting to the origin in $\operatorname{Spec} R$.

Now on X^{I_1}, the black dot is the singularity $uv = f_1$, whereas the yellow dot is $uv = f_2f_3f_4$. On X^{I_2}, the latter singularity splits into $uv = f_2$ (middle black dot) and $uv = f_3f_4$ (yellow). The left hand black dot is still $uv = f_1$. On $X^{\mathcal{F}}$, the black dots are (reading left to right) $uv = f_1$, $uv = f_2$, $uv = f_3$ and $uv = f_4$. All yellow dots correspond to non-$\mathbb{Q}$-factorial singularities, and all black dots correspond to isolated $\mathbb{Q}$-factorial singularities, possibly smooth. The yellow dots with squiggles through them are non-isolated singularities. In this example, so that we can draw a picture, we have assumed that $(f_2) = (f_3)$, and $(f_3) \neq (f_4)$, so that $uv = f_2f_3f_4$ is non-isolated, but $uv = f_3f_4$ is isolated.

The red curves are the $\mathbb{P}^1$'s which map to the origin in $\operatorname{Spec} R$. The right hand curve in X^{I_2} moves in a family (represented as blue lines) and φ_2 contracts the whole family; consequently φ_2 contracts a divisor. However both φ_1 and φ_3 contract a single curve but no divisor and so are flopping contractions. Note that $\operatorname{Spec} R$ and X^{I_1} have canonical singularities, but X^{I_2} and $X^{\mathcal{F}}$ have only terminal singularities.

The precise form of the quiver relies on 5.21, but this can be ignored for now. Note that the geometric picture will change depending both on the choice of polynomials, and their ordering. Blowing up in a different order can change which curves move in families, and also normal bundles of the curves. Of course, this depends on the choice of the polynomials too. On the algebraic side, if we change the polynomials or their order then the number of loops change, as do the relations. Changing the ordering corresponds to mutation (see 6.13 later).

On each level i the geometric space X^{I_i} is derived equivalent to the corresponding algebra by 4.2 . The top space has only $\mathbb{Q}$-factorial terminal singularities, hence the top algebra is a MMA.

Remark 4.7. We believe that with the precise conditions on f as in 4.2, the result is true in the non-local case. However, this (and also the more general non-local situation) is quite a delicate problem, even when crepant resolutions exist. For example, in the case $R = \mathbb{C}[u,v,x,y]/uv - xy(y-2)$, define the modules $M := (u,x)$, $L_1 := (u, x(y-2))$ and $L_2 := (u, xy)$. Then $M := R \oplus M_1$ and $L := R \oplus L_1 \oplus L_2$ both give NCCRs (we can check this complete locally), but since $\operatorname{add} M = \operatorname{add} L$ (again this can be checked complete locally by [IW10, 4.11]), they are Morita equivalent. By [V04b], inspecting their moduli we obtain a derived equivalence between the NCCRs and all crepant resolutions of $\operatorname{Spec} R$.

Hence our picture in 4.6 becomes



where φ_1 is obtained by blowing up the ideal (u,x). This relies on the fact that we a priori know that we have a NCCR, because we can check this (complete) locally. In a similar case $uv = (x^2+y^3)y(y-2)$ (where no NCCR exists), we cannot play this trick, since we do not know if MMAs can be detected locally, and also since completion destroys $\mathbb{Q}$-factoriality we cannot reduce to the complete local setting. If instead we start with the space with two $\mathbb{P}^1$'s and try to obtain a direct proof of the derived equivalence between the space and the two algebras (as in 4.2), things are also complicated by the fact that it is unclear which of the two is the 'natural' algebra to aim for.

5. Classification of MM modules on cA_n singularities

In this section we give a proof of a strengthened version of 4.4, using only commutative algebra. As stated in the introduction, this is important since given a 3-fold singularity R, often the explicit geometry of a $\mathbb{Q}$-factorial terminalization of $\operatorname{Spec} R$ is not known, so in such situations creating techniques to be able to argue directly on the base R is desirable.

We keep the notation as in §4.1, and work over a field k. When k is algebraically closed, we denote it $\Bbbk$. Set

$$S := k[[x,y]] \quad \text{and} \quad R := S[[u,v]]/(f(x,y)-uv).$$

We let $f = f_1 \dots f_n$ be a factorization into prime elements of S. Recall for any subset $I \subseteq \{1,\dots,n\}$ we denote

$$f_I := \prod_{i \in I} f_i \quad \text{and} \quad T_I := (u, f_I)$$

which is an ideal of R. For a collection of subsets $\emptyset \subsetneq I_1 \subsetneq I_2 \subsetneq \dots \subsetneq I_m \subsetneq \{1,2,\dots,n\}$, we say that $\mathcal{F} = (I_1,\dots,I_m)$ is a *flag in the set* $\{1,2,\dots,n\}$. We say that the flag $\mathcal{F}$ is *maximal* if $n = m+1$. Given a flag $\mathcal{F} = (I_1,\dots,I_m)$, we define

$$T^{\mathcal{F}} := R \oplus \left(\bigoplus_{j=1}^{m} T_{I_j} \right).$$

Again, as in §4.1, we can (and do) identify maximal flags with elements of the symmetric group, so for $\omega \in \mathfrak{S}_n$, we let

$$T^{\omega} := \bigoplus_{j=1}^{n} T_{\{\omega(1),\dots,\omega(j)\}}.$$

5.1. **Indecomposable modifying modules on cA_n singularities.** The first main result 5.1 classifies the indecomposable modifying R-modules which are CM, generalizing [DH10, 4.3] and [V04a, A.1] to cover non-isolated singularities.

Theorem 5.1. *When $k = \Bbbk$, $\{T_I \mid I \subseteq \{1, \dots, n\}\}$ are all the indecomposable modifying CM R-modules.*

The proof of 5.1 requires some preparation. Let us first recall the following famous Nagata's Theorem.

Proposition 5.2. *(Nagata's Theorem)* [R98, 1.2.4] *Let T be a domain and S a multiplicative set of T. Then there exists a surjective group homomorphism $\mathrm{Cl}(T) \to \mathrm{Cl}(T_S)$, $I \to I_S$ whose kernel is generated by height one prime ideals I with $I \cap S \neq \emptyset$.*

We also need the following general result to determine the precise structure of the class group. We denote by $\mathrm{Ch}(R)$ the Chow group of R.

Proposition 5.3. [R98, Section 1.2] *We have an injective homomorphism $\mathrm{Cl}(R) \to \mathrm{Ch}(R)$ sending I to $\sum_{\mathrm{ht}\mathfrak{p}=1} \mathrm{length}_{A_\mathfrak{p}}((R/I)_\mathfrak{p})[\mathfrak{p}]$.*

These give the following description of the class group of R. Note that part (3) in the following lemma is used implicitly in [DH10, 4.3], and its proof was communicated to us by Hailong Dao. We record it here for completeness.

Lemma 5.4. (1) *$\mathrm{Cl}(R)$ is generated by the ideals (u, f_i) for $1 \leq i \leq n$.*
(2) *Suppose that $g, h \in k[[x, y]]$ such that gh divides f. Then $[(u, g)] + [(u, h)] = [(u, gh)]$.*
(3) *Every CM element in $\mathrm{Cl}(R)$ has the form (u, f_I) for some $I \subseteq \{1, \dots, n\}$.*

Proof. (1) Applying Nagata's Theorem for the multiplicative set S generated by u, we have a surjective group homomorphism $\mathrm{Cl}(R) \to \mathrm{Cl}(R_S)$ whose kernel is generated by height one prime ideals I with $I \cap S \neq \emptyset$. Since

$$R_S \simeq k[[x, y, u, u^{-1}, v]]/(v + u^{-1} f) \simeq k[[x, y, u, u^{-1}]]$$

is a factorial domain, we have $\mathrm{Cl}(R_S)$ is trivial. Thus $\mathrm{Cl}(R)$ is generated by height one prime ideals intersecting with S.

Let I be a height one prime ideal of R intersecting with S. Then $u^i \in I$ for some $i > 0$, so we have $u \in I$ since I is a prime ideal. Now consider the surjective map

$$R = k[[x, y, u, v]]/(f - uv) \to R/(u) = k[[x, y, u, v]]/(f, u) = k[[x, y, v]]/(f).$$

Since I is a height one prime ideal of R, the ideal $I/(u)$ is a minimal prime ideal of $k[[x, y, v]]/(f)$. Thus $I/(u)$ has to be $(f_i)/(f)$ for some i, and so we have $I = (u, f_i)$.
(2) Since gh divides f, by considering the appropriate matrix factorization, we see that $(u, gh) \in \mathrm{CM}\, R$ and (u, gh) has rank one. It follows that $(u, gh) \in \mathrm{Cl}(R)$ and hence by 5.3 we only have to show that

$$\mathrm{length}_{A_\mathfrak{p}}((R/(u,g))_\mathfrak{p}) + \mathrm{length}_{A_\mathfrak{p}}((R/(u,h))_\mathfrak{p}) = \mathrm{length}_{A_\mathfrak{p}}((R/(u,gh))_\mathfrak{p})$$

for all height one primes $\mathfrak{p}$. We know that $R/(u, g) = k[[x, y, v]]/(g)$, $R/(u, h) = k[[x, y, v]]/(h)$ and $R/(u, gh) = k[[x, y, v]]/(gh)$. Since we have an exact sequence

$$0 \to k[[x, y, v]]/(g) \xrightarrow{h} k[[x, y, v]]/(gh) \to k[[x, y, v]]/(h) \to 0,$$

the desired equality holds by localizing at $\mathfrak{p}$.
(3) Let $I = (x, y, u - v)$ be a system of parameters of R, and let $\mathfrak{n}$ denote the maximal ideal of R. Suppose that M is a CM member of the class group. Since $R/I \simeq k[u]/(u^2)$, we have $\mathrm{length}(R/I) = 2$. Further

$$\mathrm{length}(M/\mathfrak{n}M) \leq \mathrm{length}(M/IM).$$

Now by [BH, 4.7.11(3)]

$$\mathrm{length}(M/IM) = \mathrm{length}(R/I) \times \mathrm{rank}(M) = 2,$$

and so the minimal number of generators of M is less than or equal to two. If it is one then there is nothing to prove, so assume that it is two. Pulling M across Knöerrer periodicity (say to M_1), then the minimal number of generators of M_1 is one. This means that there must be a matrix factorization

$$k[[x,y]] \xrightarrow{g} k[[x,y]] \xrightarrow{h} k[[x,y]] \to M_1 \to 0$$

where h must divide f, say $h = f_I$. Pulling back across Knöerrer periodicity, we see that M must have the form (u, f_I). $\square$

Moreover, we require the following lemma:

Lemma 5.5. *Suppose that $k = \Bbbk$. Then for any modifying R-module M, we can choose a hyperplane section t satisfying the following two conditions.*
(1) $R/(t)$ is an A_m singularity, where m is the degree of the lowest term of f minus one.
(2) t acts on $E := \operatorname{Ext}^1_R(M,M)$ as a non-zerodivisor.

Proof. Since R is a cA_m singularity (see e.g [BIKR, 6.1(e)]), a generic hyperplane section t satisfies condition (1). If t acts on E as a zero divisor, then t is contained in an associated prime ideal of E. But $\operatorname{depth}_R E > 0$ by [IW10, 2.6], so any associated prime ideal of E is necessarily non-maximal. Since E has only finitely many associated prime ideals, we can find a hyperplane t which is not contained in any associated prime ideal of E, and furthermore satisfies (1). $\square$

Now we are ready to prove 5.1.

Proof. Suppose that $M \in \operatorname{CM} R$ satisfies $\operatorname{End}_R(M) \in \operatorname{CM} R$, then certainly we have $\operatorname{depth}_R \operatorname{Ext}^1_R(M,M) > 0$ by 2.9. We first claim that M has rank one, so suppose on the contrary that $\operatorname{rk} M > 1$.

By 5.5 we can pick $t \in R$ such that $R_1 := R/(t) \cong \Bbbk[[x,y,z]]/(x^2+y^2+z^m)$, and t acts on $\operatorname{Ext}^1_R(M,M)$ as a non-zerodivisor. Denote $\Lambda := \operatorname{End}_R(M)$, then applying $\operatorname{Hom}_R(M,-)$ to the exact sequence

$$0 \to M \xrightarrow{t} M \to M/tM \to 0$$

yields

$$0 \to \Lambda \xrightarrow{t} \Lambda \to \operatorname{Hom}_R(M, M/tM) \to \operatorname{Ext}^1_R(M,M) \xrightarrow{t} \operatorname{Ext}^1_R(M,M)$$

Since t acts on $\operatorname{Ext}^1_R(M,M)$ as a non-zerodivisor, we have

$$\Lambda/t\Lambda \simeq \operatorname{Hom}_R(M, M/tM) \simeq \operatorname{End}_R(M/tM) = \operatorname{End}_{R_1}(M/tM).$$

Since R_1 is a simple surface singularity of type A_{m-1}, it is well-known (e.g. [Yos90]) that all indecomposable CM R_1-modules have rank one. Since $\operatorname{rk}_{R_1}(M/tM) = \operatorname{rk}_R(M) > 1$, we have that M/tM is decomposable. Thus $\Lambda/t\Lambda \simeq \operatorname{End}_{R_1}(M/tM)$ has a non-trivial idempotent. Since R is complete, $\Lambda = \operatorname{End}_R(M)$ also has a non-trivial idempotent by the lifting idempotents property, a contradiction to indecomposability of M. Hence $\operatorname{rk} M = 1$, so we can regard M as an element of $\operatorname{Cl}(R)$. By 5.4(3), we have the assertion. $\square$

5.2. MM and modifying modules on cA_n singularities.

In 5.1 we classified indecomposable modifying R-modules, and we now proceed to prove 5.16, where we classify all basic modifying R-modules, and all basic maximal modifying R-modules. Although these statements are for the three dimensional ring R, to prove them we need to pass to the one-dimensional ring

$$R^\flat := S/(f).$$

There is a triangle equivalence

$$K : \underline{\operatorname{CM}}\, R^\flat \to \underline{\operatorname{CM}}\, R$$

called *Knöerrer periodicity* (valid for any field k by [S89, 3.1]) which is defined as follows [Yos90]: For any $X \in \operatorname{CM} R^\flat$, we have a free resolution

$$R^{\flat \oplus n} \xrightarrow{B} R^{\flat \oplus n} \xrightarrow{A} R^{\flat \oplus n} \to X \to 0$$

where A and B are $n \times n$ matrices over S satisfying $AB = BA = f 1_{S^n}$. Then KX is defined by the following exact sequence:

$$R^{\oplus 2n} \xrightarrow{\left(\begin{smallmatrix} B & -u \\ -v & A \end{smallmatrix}\right)} R^{\oplus 2n} \xrightarrow{\left(\begin{smallmatrix} A & u \\ v & B \end{smallmatrix}\right)} R^{\oplus 2n} \to KX \to 0.$$

We use the following terminology, which will be justified by Knörrer periodicity.

Definition 5.6. *(1) We say that $X \in \operatorname{CM} R^\flat$ is a* $\flat$-modifying module *if*

$$\operatorname{depth}_{R^\flat} \operatorname{Ext}^1_{R^\flat}(M,M) > 0.$$

(2) We say that $X \in \operatorname{CM} R^\flat$ is a maximal $\flat$-modifying (=MM$^\flat$) module *if it is a $\flat$-modifying module, and further if $X \oplus Y$ is a $\flat$-modifying module for some $Y \in \operatorname{CM} R^\flat$, then $Y \in \operatorname{add} X$.*
(3) We call $M \in \operatorname{CM} R^\flat$ a CT$^\flat$ module *if*

$$\operatorname{add} M = \{X \in \operatorname{CM} R^\flat \mid \operatorname{depth} \operatorname{Ext}^1_{R^\flat}(M,X) > 0\}.$$

Note that since $R^\flat$ has dimension one, $\flat$-modifying $R^\flat$-modules do not correspond to modifying $R^\flat$-modules. However, we will see in 5.11 that $\flat$-modifying $R^\flat$-modules correspond to modifying R-modules, and that CT$^\flat$ $R^\flat$-modules correspond to CT R-modules.

For any subset $I \subseteq \{1,\dots,n\}$, and any flag $\mathcal{F} = (I_1,\dots,I_m)$ in the set $\{1,2,\dots,n\}$, we define

$$S_I := S/(f_I) \cong R^\flat/(f_I) \quad \text{and} \quad S^{\mathcal{F}} := R^\flat \oplus \left(\bigoplus_{j=1}^{m} S_{I_j} \right).$$

Again we identify maximal flags with elements of the symmetric group, and so for $\omega \in \mathfrak{S}_n$ we set $S^\omega := \bigoplus_{j=1}^n S_{\{\omega(1),\dots,\omega(j)\}}$. The following is immediate:

Lemma 5.7. *Let $\mathcal{F}$ be a flag in the set $\{1,\dots,n\}$. Then the Knöerrer periodicity functor sends $S^{\mathcal{F}}$ (respectively, S^ω) to $T^{\mathcal{F}}$ (respectively, T^ω).*

The basic strategy is to prove the main theorem (5.16) by reduction to the one-dimensional case. It is easy to show that the property cluster–tilting is invariant under any triangle equivalence $\underline{\operatorname{CM}} R_1 \approx \underline{\operatorname{CM}} R_2$ where R_1 and R_2 are commutative Gorenstein rings, however replacing questions involving Ext vanishing with questions involving Ext having positive depth is a little more subtle.

First, we note that for any commutative ring R, the categories $\underline{\operatorname{CM}} R$ and $\operatorname{CM} R$ are *R-linear*, i.e. R acts on each morphism set so that the composition is R-bilinear.

Lemma 5.8. *Let R_i $(i=1,2)$ be commutative rings and let $\mathcal{C}_i$ be R_i-linear categories $(i=1,2)$. Suppose that $F : \mathcal{C}_1 \to \mathcal{C}_2$ is an equivalence of additive categories. Then for all $X, Y \in \mathcal{C}_1$, the action of R_1 on $\operatorname{Hom}_{\mathcal{C}_1}(X,Y)$ and the action of R_2 on $\operatorname{Hom}_{\mathcal{C}_2}(FX,FY)$ commute under the isomorphism $F : \operatorname{Hom}_{\mathcal{C}_1}(X,Y) \simeq \operatorname{Hom}_{\mathcal{C}_2}(FX,FY)$.*

Proof. Since $\mathcal{C}_1$ is R_1-linear, for $r \in R_1$ and $\alpha \in \operatorname{Hom}_{\mathcal{C}_1}(X,Y)$, we have

$$r \cdot \alpha = (X \xrightarrow{r1_X} X \xrightarrow{\alpha} Y) = (X \xrightarrow{\alpha} Y \xrightarrow{r1_Y} Y). \tag{5.F}$$

Similarly, since $\mathcal{C}_2$ is R_2-linear, for $r' \in R_2$ and $\beta \in \operatorname{Hom}_{\mathcal{C}_2}(X',Y')$, we have

$$r' \cdot \beta = (X' \xrightarrow{r'1_{X'}} X' \xrightarrow{\beta} Y') = (X' \xrightarrow{\beta} Y' \xrightarrow{r'1_{Y'}} Y'). \tag{5.G}$$

We denote by $\star$ the action of R_1 on $\operatorname{Hom}_{\mathcal{C}_2}(FX,FY)$ induced by the isomorphism $F : \operatorname{Hom}_{\mathcal{C}_1}(X,Y) \simeq \operatorname{Hom}_{\mathcal{C}_2}(FX,FY)$. Then for $r \in R_1$ and $\beta \in \operatorname{Hom}_{\mathcal{C}_2}(FX,FY)$

$$r \star \beta = F(r \cdot F^{-1}(\beta)) \overset{(5.F)}{=} F(X \xrightarrow{r1_X} X \xrightarrow{F^{-1}(\beta)} Y) = (FX \xrightarrow{F(r1_X)} FX \xrightarrow{\beta} FY). \tag{5.H}$$

Now for all $r \in R_1$, $r' \in R_2$ and $\beta \in \operatorname{Hom}_{\mathcal{C}_2}(FX,FY)$

$$\begin{aligned} r' \cdot (r \star \beta) &\overset{(5.H)}{=} r' \cdot (FX \xrightarrow{F(r1_X)} FX \xrightarrow{\beta} FY) \overset{(5.G)}{=} (FX \xrightarrow{F(r1_X)} FX \xrightarrow{\beta} FY \xrightarrow{r'1_Y} FY) \\ &\overset{(5.H)}{=} r \star (FX \xrightarrow{\beta} FY \xrightarrow{r'1_Y} FY) \overset{(5.G)}{=} r \star (r' \cdot \beta), \end{aligned}$$

as required. ☐

Recall from §2.3 that if $X \in \operatorname{mod} R$, we denote $\mathsf{fl}_R X$ to be the largest finite length R-submodule of X.

Lemma 5.9. *Suppose that k is a field and let $(R_1,\mathfrak{m})$ and $(R_2,\mathfrak{n})$ be commutative noetherian local finitely generated k-algebras. Let M be an (R_1,R_2)-bimodule and assume*
(i) $(\lambda 1_{R_1})\cdot m = m\cdot(\lambda 1_{R_2})$ *for all* $\lambda\in k$.
(ii) *M is finitely generated as an R_1-module and as an R_2-module.*
Then $\mathsf{fl}_{R_1} M = \mathsf{fl}_{R_2} M$. *In particular* $\operatorname{depth}_{R_1} M > 0$ *if and only if* $\operatorname{depth}_{R_2} M > 0$.

Proof. $\mathsf{fl}_{R_2} M = \{x\in M \mid \exists\, r\in\mathbb{N} \text{ with } x\mathfrak{n}^r = 0\}$ is a sub R_1-module of M. Since $M\in\operatorname{mod} R_2$ is finitely generated and $R_2/\mathfrak{n}$ is a finite dimensional k-space, we have that $\dim_k \mathsf{fl}_{R_2} M < \infty$. By (i) the action of k viewed inside R_1 and the action of k viewed inside R_2 coincide, so $\mathsf{fl}_{R_2} M$ is a module for the k-algebra R_1 for which $\dim_k \mathsf{fl}_{R_2} M < \infty$. Certainly this implies that $\mathsf{fl}_{R_2} M$ must have finite length as an R_1-module, and so $\mathsf{fl}_{R_2} M \subseteq \mathsf{fl}_{R_1} M$. We have the opposite inclusion similarly. The last statement holds since for any commutative local noetherian ring R and any $M\in\operatorname{mod} R$, $\operatorname{depth}_R M > 0 \iff \mathsf{fl}_R M = 0$. ☐

We now apply the above to Knörrer periodicity. The following is now immediate:

Proposition 5.10. *With notation as above, for all $M,N\in\operatorname{CM} R^\flat$ we have*
(1) $\operatorname{depth}_{R^\flat} \underline{\operatorname{Hom}}_{R^\flat}(M,N) > 0 \iff \operatorname{depth}_R \underline{\operatorname{Hom}}_R(KM,KN) > 0$.
(2) $\operatorname{depth}_{R^\flat} \operatorname{Ext}^1_{R^\flat}(M,N) > 0 \iff \operatorname{depth}_R \operatorname{Ext}^1_R(KM,KN) > 0$.

Proof. (1) $\underline{\operatorname{Hom}}_{R^\flat}(M,N)\cong\underline{\operatorname{Hom}}_R(KM,KN)$ is a $(R^\flat,R)$-bimodule by 5.8. By construction the Knörrer functor K is k-linear, so the condition in 5.9(i) holds. Since the condition 5.9(ii) is clear, the result follows by 5.9.
(2) We only have to apply (1) by replacing N by $N[1]$ and use the fact that K is a triangle functor. ☐

Corollary 5.11. *Suppose that $M\in\operatorname{CM} R^\flat$ and $N\in\operatorname{CM} R$ are generators such that $KM\cong N$. Then*
(1) *M is a $\flat$-modifying $R^\flat$-module $\iff$ N is modifying R-module.*
(2) *M is a MM$^\flat$ $R^\flat$-module $\iff$ N is a MM R-module.*
(3) *M is a CT$^\flat$ $R^\flat$-module $\iff$ N is a CT R-module.*

Proof. The assertions follow from 5.10. ☐

The following two lemmas are the key calculations from dimension one that are needed in the proof of the main theorem (5.15).

Lemma 5.12. *Let $R^\flat$ be a local hypersurface of dimension one, $M,N\in\operatorname{CM} R^\flat$. Then*
$$M^*\otimes_{R^\flat} N\in\operatorname{CM} R^\flat \iff \operatorname{depth}_{R^\flat}\operatorname{Ext}^1_{R^\flat}(M,N)>0.$$

Proof. Using the exact sequence
$$0\to\operatorname{Ext}^1_{R^\flat}(\operatorname{Tr} N,M^*)\xrightarrow{f} N\otimes_{R^\flat} M^*\to\operatorname{Hom}_{R^\flat}(N^*,M^*)\to\operatorname{Ext}^2_{R^\flat}(\operatorname{Tr} N,M^*)\to 0$$
certainly $\operatorname{depth}_{R^\flat}\operatorname{Cok} f>0$. Hence $M^*\otimes_{R^\flat} N\in\operatorname{CM} R^\flat \iff \operatorname{depth}_{R^\flat}\operatorname{Ext}^1_{R^\flat}(\operatorname{Tr} N,M^*)>0$. But
$$\operatorname{Ext}^1_{R^\flat}(\operatorname{Tr} N,M^*)=\underline{\operatorname{Hom}}_{R^\flat}(\Omega\operatorname{Tr} N,M^*)=\underline{\operatorname{Hom}}_{R^\flat}((\Omega N)^*,M^*)=\underline{\operatorname{Hom}}_{R^\flat}(M,\Omega N)$$
and since $R^\flat$ is a hypersurface $\Omega N=\Omega^{-1}N$. Thus
$$\operatorname{Ext}^1_{R^\flat}(\operatorname{Tr} N,M^*)=\underline{\operatorname{Hom}}_{R^\flat}(M,\Omega N)=\underline{\operatorname{Hom}}_{R^\flat}(M,\Omega^{-1}N)=\operatorname{Ext}^1_{R^\flat}(M,N),$$
completing the proof. ☐

Lemma 5.13. *Let I and J be subsets of $\{1,\dots,n\}$. Then*
(1) $S_I\cong S_I^*$.
(2) $S_I\otimes_{R^\flat} S_J\in\operatorname{CM} R^\flat \iff I\subseteq J$ *or* $J\subseteq I$.

Proof. (1) is an easy calculation.
(2) First, observe that $S_I\otimes_{R^\flat} S_J\cong R^\flat/(f_I)\otimes_{R^\flat} R^\flat/(f_J)\cong R^\flat/(f_I,f_J)\cong S/(f_I,f_J)$.
($\Leftarrow$) If $I\subseteq J$ then $f_J\in(f_I)$, so $(f_I,f_J)=(f_I)$ and thus $S_I\otimes_{R^\flat} S_J\cong S_I\in\operatorname{CM} R^\flat$. The case $J\subseteq I$ is identical.
($\Rightarrow$) Suppose that $S/(f_I,f_J)\in\operatorname{CM} R^\flat$. Then, as is standard in matrix factorizations, the kernel of the natural map $S\to S/(f_I,f_J)\to 0$ is a free S-module, necessarily of rank

one. But the kernel is clearly (f_I, f_J), and so in particular (f_I, f_J) must be generated by a single element, say $(f_I, f_J) = (t)$. Clearly t must divide f_I and f_J, and so (since S is a factorial domain) it must divide their highest common factor. If $I \nsubseteq J$ and $J \nsubseteq I$ then this highest common factor is strictly smaller that both f_I and f_J, hence we may write $f_I = at$ and $f_J = bt$ for some non-units a and b. But now $t \in (f_I, f_J)$, so we may write

$$t = cf_I + df_J = (ac + bd)t$$

for some $c, d \in S$. But S is a domain, so this forces $1 = ac + bd$. Since S is local, we deduce that either ac or bd is a unit. Either way this is impossible, since a and b are not units. Hence either $I \subseteq J$ or $J \subseteq I$. ☐

Corollary 5.14. (1) $\operatorname{depth}_{R^\flat} \operatorname{Ext}^1_{R^\flat}(S_I, S_J) > 0 \iff I \subseteq J$ *or* $J \subseteq I$.
(2) *For all flags* $\mathcal{F} = (I_1, \dots, I_m)$, $\operatorname{depth}_{R^\flat} \operatorname{Ext}^1_{R^\flat}(S^{\mathcal{F}}, S^{\mathcal{F}}) > 0$.

Proof. (1) By 5.13(1) $(S_I)^* \otimes_{R^\flat} S_J \cong S_I \otimes_{R^\flat} S_J$. Hence the result follows by combining 5.13(2) and 5.12.
(2) Follows by repeated application of (1). ☐

Recall that we say that an R-module is *basic* if it is isomorphic to a finite direct sum of mutually non-isomorphic indecomposable R-modules.

Theorem 5.15. *Suppose that* $k = \mathbb{k}$. *Then for each flag* $\mathcal{F}$ *in the set* $\{1, \dots, n\}$,
(1) $S^{\mathcal{F}}$ *is a basic* $\flat$*-modifying CM* $R^\flat$*-module. All basic* $\flat$*-modifying CM* $R^\flat$*-modules arise in this way.*
(2) *Under the assignment in* (1)*, maximal flags give maximal* $\flat$*-modifying CM* $R^\flat$*-modules, and further all basic maximal* $\flat$*-modifying CM* $R^\flat$*-modules arise from maximal flags.*

Proof. (1) We know that $S^{\mathcal{F}}$ is $\flat$-modifying by 5.14(2). Conversely, let Z be a basic $\flat$-modifying $R^\flat$-module. By combining 5.1, 5.7 and 5.11, Z is a direct sum of some S_I. Now Z has the form $S^{\mathcal{F}}$ for some $\mathcal{F}$ by repeated use of 5.14(1).

For maximality, let $\mathcal{F}$ be a maximal flag. Suppose that $S^{\mathcal{F}} \oplus Y$ is $\flat$-modifying for some $Y \in \operatorname{CM} R^\flat$. Since Y itself must be $\flat$-modifying, by above $Y = S^{\mathcal{G}}$ for some flag $\mathcal{G}$. That $Y \in \operatorname{add} S^{\mathcal{F}}$ now follows by repeated use of 5.14(1). Conversely, let Z be a basic maximal $\flat$-modifying $R^\flat$-module. By above, $Z = S^{\mathcal{F}}$ for some flag $\mathcal{F}$. If $\mathcal{F}$ is not maximal we can refine $\mathcal{F}$ to a new flag $\mathcal{G}$ with more terms. This would mean that $S^{\mathcal{G}}$ is a $\flat$-modifying $R^\flat$-module, containing $Z = S^{\mathcal{F}}$ as a summand, such that $S^{\mathcal{G}}$ has a summand which does not belong to $\operatorname{add} Z$. This contradicts maximality of Z, hence $\mathcal{F}$ must have been maximal. ☐

The following is the main result of this subsection.

Corollary 5.16. *Suppose that* $k = \mathbb{k}$. *Then for each flag* $\mathcal{F}$ *in the set* $\{1, \dots, n\}$,
(1) $T^{\mathcal{F}}$ *is a basic modifying CM* R*-module. All basic modifying CM* R*-modules arise in this way.*
(2) *Under the assignment in* (1)*, maximal flags give maximal modifying CM* R*-modules, and further all basic maximal modifying CM* R*-modules arise from maximal flags.*

Proof. This is immediate from 5.15, 5.11 and 5.7. ☐

This leads to the following strengthened version of 4.4, using completely different methods.

Corollary 5.17. *Suppose that* $k = \mathbb{k}$ *and let* $R = \mathbb{k}[[u, v, x, y]]/(uv - f_1 \dots f_n)$. *Then the MM* R*-module generators are precisely* T^ω, *where* $\omega \in \mathfrak{S}_n$.

Proof. This is just a reformulation of 5.16(2), since we can identify maximal flags with members of the symmetric group. ☐

Corollary 5.18. *Suppose that* $k = \mathbb{k}$. *Grouping together terms in the decomposition of* f, *write* $f = f_1^{a_1} \dots f_t^{a_t}$ *for some* distinct *prime elements* f_i, *and some* $a_i \in \mathbb{N}$. *Then* R *has precisely* $\frac{(a_1+\dots+a_t)!}{a_1!\dots a_t!}$ *basic MM generators.*

Proof. All basic MM generators have the form T^ω for some $\omega \in \mathfrak{S}_{a_1+\dots+a_t}$ by 5.16(2). Accounting for the repetitions, there are precisely $\frac{(a_1+\dots+a_t)!}{a_1!\dots a_t!}$ of these. ☐

We will describe these MM generators pictorially in §6.

5.3. **CT modules on cA_n singularities.** Given the existence of MM generators, it is natural to ask under what circumstances they are CT. The following is a classification of the CT modules, strengthening 4.4 and generalizing both [BIKR, 1.5] and [DH10, 4.2] to non-isolated singularities.

Theorem 5.19. *Suppose that $k = \Bbbk$. Then the following conditions are equivalent.*
(1) *$R^\flat$ has a $CT^\flat$ module.*
(2) *R has a CT module.*
(3) *$f_i \notin \mathfrak{m}^2$ for all $1 \le i \le n$.*
In this case, all basic $CT^\flat$ $R^\flat$-modules are of the form S^ω for some $\omega \in \mathfrak{S}_n$, and all basic CT R-modules are of the form T^ω for some $\omega \in \mathfrak{S}_n$.

We prove this on p.27. The trick is not to compute the global dimension of an arbitrary MMA (which is hard), but a well-chosen one. As soon as one MMA has finite global dimension, they all do [IW10, 5.14].

We first calculate the quiver of $\mathrm{End}_R(T^{\mathcal{F}})$, which requires the following easy lemma.

Lemma 5.20. *Let $R = k[[u,v,x,y]]/(uv-f)$, and consider two modifying R-modules T_I and T_J such that either $I \subset J$ or $J \subset I$. Then*

$$\mathrm{Hom}_R(T_I, T_J) \cong \begin{cases} (u, \frac{f_J}{f_I}) & \text{if } I \subset J \\ (u, \frac{f_J f}{f_I}) & \text{if } J \subset I \end{cases}$$

Proof. Since R is normal we have $\mathrm{Hom}_R(T_I, T_J) \cong ((u, f_I)^* \otimes_R (u, f_J))^{**}$. Now if $I \subset J$ then f_I divides f_J, so write $f_J = \frac{f_J}{f_I} f_I$. But now by 5.3(2)

$$[(u, \tfrac{f_J}{f_I})] + [(u, f_I)] = [(u, \tfrac{f_J}{f_I} f_I)] = [(u, f_J)],$$

from which we deduce that $((u, f_I)^* \otimes_R (u, f_J))^{**} \cong (u, \frac{f_J}{f_I})$. For the second statement, suppose that $J \subset I$, then f_J divides f_I, so

$$[(u, \tfrac{f_I}{f_J})] + [(u, f_J)] = [(u, f_I)]$$

shows that $((u, f_I)^* \otimes_R (u, f_J))^{**} \cong (u, \frac{f_I}{f_J})^*$. But since $[(u, \frac{f_I}{f_J})] + [(u, \frac{f_J f}{f_I})] = [(u, f)] = 0$, it follows that $(u, \frac{f_I}{f_J})^* \cong (u, \frac{f_J f}{f_I})$, as required. □

To ease notation in the following quiver, for a given flag $\mathcal{F} = (I_1, \dots, I_m)$ we denote $g_1 := f_{I_1}$, set $g_j := \frac{f_{I_j}}{f_{I_{j-1}}}$ for all $2 \le j \le m$, and $g_{m+1} := \frac{f}{f_{I_m}}$.

Corollary 5.21. *Given a flag $\mathcal{F} = (I_1, \dots, I_m)$, with notation as above the quiver of $\mathrm{End}_R(T^{\mathcal{F}})$ is as follows:*



$m{\ge}2$



$m{=}1$

together with the possible addition of some loops, given by the following rules:

- *Consider vertex R. If $(g_1, g_{m+1}) = (x, y)$ in the ring $k[[x,y]]$, add no loops at vertex R. Hence suppose $(g_1, g_{m+1}) \subsetneq (x, y)$. If there exists $t \in (x, y)$ such that $(g_1, g_{m+1}, t) = (x, y)$, add a loop labelled t at vertex R. If there exists no such t, add two loops labelled x and y at vertex R.*
- *Consider vertex T_{I_i}. If $(g_i, g_{i+1}) = (x, y)$ in the ring $k[[x,y]]$, add no loops at vertex T_{I_i}. Hence suppose $(g_i, g_{i+1}) \subsetneq (x, y)$. If there exists $t \in (x, y)$ such that $(g_i, g_{i+1}, t) = (x, y)$, add a loop labelled t at vertex T_{I_i}. If there exists no such t, add two loops labelled x and y at vertex T_{I_i}.*

Proof. (1) Since $\mathrm{Hom}_R(R,R)\cong \mathrm{Hom}_R(T_{I_i},T_{I_i})\cong R$, we first must verify that at each vertex we can see the elements u,v,x,y as cycles at that vertex. Certainly u is there (as it is the path followed anticlockwise around the circle), and certainly v is there (being the path followed clockwise around the circle). It is possible to see cycles x and y at every vertex by the rules for loops.
(2) Since $\mathrm{Hom}_R(R,T_{I_i})\cong T_{I_i}$, we must verify that we can see the generators of T_{I_i} as paths from vertex R to vertex T_{I_i}. But this is clear, as $\prod_{j=1}^{i} g_j$ is the clockwise path, and u is the anticlockwise path.
(3) As a module over the centre, the paths from vertex T_{I_i} to vertex R are generated by two paths from vertex T_{I_i} to R, namely the one clockwise, which is $\frac{f}{uf_I}$, and the one anticlockwise, which is inclusion. Since $\mathrm{Hom}_R(T_{I_i},R)\cong (u,\frac{f}{f_I})$ (see proof of 5.20), clearly this is isomorphic to paths from T_{I_i} to R.
(4) The argument for paths from T_{I_i} to T_{I_j} is identical to the argument in (3), using the isomorphism in 5.20. ☐

Remark 5.22. We remark that there are at least two other methods for computing the quiver of $\mathrm{End}_R(T^{\mathcal{F}})$. One way would be to use reconstruction on 1-dimensional fibres à la $\mathrm{GL}(2,\mathbb{C})$ McKay Correspondence [W10]. Another is to compute the quiver of $\mathrm{End}_R(T^{\mathcal{F}})$ in the one-dimensional setting, as in [BIKR, 4.10], and then use Knörrer periodicity. This last method only gives some of the quiver, so more work would be needed.

We will also need the following two easy lemmas.

Lemma 5.23. *Suppose that R is a complete local 3-CY$^-$ normal domain, and let M be a basic maximal modifying R-module. If in the quiver of* $\mathrm{End}_R(M)$ *there are no loops, then* $\mathrm{gl.dim}\,\mathrm{End}_R(M)=3$. *In particular, M is CT.*

Proof. Set $\Lambda:=\mathrm{End}_R(M)$, and suppose that e_i is the idempotent corresponding to a vertex i. Denote $\Lambda_i:=\Lambda/\Lambda(1-e_i)\Lambda$. Since Λ is a MMA, it is easy to check (exactly as in the argument at the start of the proof of [IW10, 6.12]) that $\mathrm{proj.dim}_\Lambda\Lambda_i\le 3$ for all i. But since there are no loops in the quiver, Λ_i is just the simple at the vertex i. Notice that $\mathrm{gl.dim}\,\Lambda$ is the supremum of the projective dimensions of the simple Λ-modules (e.g. [IR08, 2.2]), hence $\mathrm{gl.dim}\,\Lambda\le 3$. A global dimension of strictly less than three would contradict the depth lemma. The final statement is just [IW10, 5.5]. ☐

Proposition 5.24. *Suppose that $M^\flat\in \mathrm{CM}\,R^\flat$ and $M\in\mathrm{CM}\,R$ are generators such that $KM^\flat\cong M$. Assume that $M^\flat$ is a $\flat$-modifying $R^\flat$-module (equivalently, by 5.11(1), that M is a modifying R-module). Then the following are equivalent.*
(1) *$M^\flat$ is a CT$^\flat$ $R^\flat$-module.*
(2) $\mathrm{gl.dim}\,\mathrm{End}_{R^\flat}(M^\flat)\le 3$.
(3) *For all $X\in\mathrm{CM}\,R^\flat$, if $M_0^\flat\xrightarrow{f}X$ is some right* $\mathrm{add}\,M^\flat$*-approximation of X in* $\mathrm{CM}\,R^\flat$*, then* $\mathrm{Ker}\,f\in\mathrm{add}\,M^\flat$.
(4) *For all $X\in\mathrm{CM}\,R^\flat$, if $M_0^\flat\xrightarrow{f}X$ is some right* $\mathrm{add}\,M^\flat$*-approximation of X in* $\underline{\mathrm{CM}}\,R^\flat$*, then* $\mathrm{cone}(f)[-1]\in\mathrm{add}\,M^\flat$.
(5) *M is a CT R-module.*
(6) $\mathrm{gl.dim}\,\mathrm{End}_R(M)\le 3$.
(7) *For all $X\in\mathrm{CM}\,R$, if $M_0\xrightarrow{f}X$ is some right* $\mathrm{add}\,M$*-approximation of X in* $\mathrm{CM}\,R$*, then* $\mathrm{Ker}\,f\in\mathrm{add}\,M$.
(8) *For all $X\in\mathrm{CM}\,R$, if $M_0\xrightarrow{f}X$ is some right* $\mathrm{add}\,M$*-approximation of X in* $\underline{\mathrm{CM}}\,R$*, then* $\mathrm{cone}(f)[-1]\in\mathrm{add}\,M$.

Proof. (1)⇔(5) This follows from 5.11(3).
(2)⇔(3) This is routine, see e.g. proof of [IW08, 2.11].
(3)⇔(4) Since short exact sequences give triangles, and triangles come from short exact sequences, this follows directly from the following easy fact: for a surjection $f:M_0^\flat\to X$, $\mathrm{Hom}_{R^\flat}(M^\flat,M_0^\flat)\to\mathrm{Hom}_{R^\flat}(M^\flat,X)$ is surjective if and only if $\mathrm{Hom}_{\underline{\mathrm{CM}}\,R^\flat}(M^\flat,M_0^\flat)\to\mathrm{Hom}_{\underline{\mathrm{CM}}\,R^\flat}(M^\flat,X)$ is surjective. For the proof of this fact, note that the "only if" part is clear, whereas the "if" part follows directly from the surjectivity of f.

(4)⇔(8) This is clear from Knörrer equivalence.
(5)⇔(6)⇔(7) These are shown in [IW10, 5.5] and [IW10, 5.13].
(7)⇔(8) is the same as the proof that (3)⇔(4). ☐

The following is a generalization of [BIKR, 4.9] to non-isolated singularities.

Lemma 5.25. (cf. [BIKR, 4.9]) *Let $R \to S$ be a surjective ring homomorphism of one-dimensional complete local rings, both of which are hypersurfaces, and let M be a $\flat$-modifying generator of S. If $R \oplus M$ is a CT$^\flat$ R-module, then M is a CT$^\flat$ S-module.*

Proof. Although the proof is similar, we give a proof for the convenience of the reader. We only have to check the condition 5.24(3) for S. Let $X \in \operatorname{CM} S$, and let $f : M_0 \to X$ be a right $\operatorname{add} M$-approximation. Since M is a generator of S, we have that f is surjective. Thus we have an exact sequence of CM S-modules

$$0 \to Y \to M_0 \xrightarrow{f} X \to 0.$$

As an R-module, X belongs to $\operatorname{CM} R$, and clearly f is a right $\operatorname{add} R \oplus M$-approximation. Since $R \oplus M$ is a CT$^\flat$ R-module, we have $Y \in \operatorname{add} R \oplus M$ by 5.24(1)⇒(3). Since Y is an S-module, we have $Y \in \operatorname{add} M$. ☐

Now we are ready to prove 5.19.

Proof. (1)⇔(2) This is clear from 5.11(3).
(2)⇒(3) We use induction on n. If $n = 1$, then R is a MM R-module by 5.1. But by assumption R has a CT module, thus by 2.11 R is a CT R-module. This implies that R is regular, hence since k is algebraically closed we can use the Jacobi criterion to deduce that $f_1 \notin \mathfrak{m}^2$. Thus the statement is true for $n = 1$, so assume that the statement is true for $n - 1$. Since there exists a CT R-module, MM generators are precisely the CT R-modules by 2.11. Hence T^ω, being a MM generator by 5.16, is CT for all $\omega \in \mathfrak{S}_n$. It follows from 5.11(3) that S^ω is a CT$^\flat$ $R^\flat$-module for all $\omega \in \mathfrak{S}_n$. Now applying 5.25 to the surjective ring homomorphism $R^\flat \to S^\omega_{n-1}$, it follows that $S^\omega_1 \oplus \cdots \oplus S^\omega_{n-1}$ is a CT$^\flat$ S^ω_{n-1}-module. By the assumption of induction, we have $f_{\omega(i)} \notin \mathfrak{m}^2$ for all $1 \le i \le n - 1$. Since ω is arbitrary, we have $f_i \notin \mathfrak{m}^2$ for all $1 \le i \le n$.
(3)⇒(1) Assume that every $f_i \notin \mathfrak{m}^2$, then we may find $g_i \in \mathfrak{m}$ such that

$$(f_i, g_i) = (g_i, f_{i+1}) = \mathfrak{m}$$

for all $1 \le i \le n - 1$. Hence consider the ring

$$R_{\text{new}} := \Bbbk[[u, v, x, y]]/(uv - f_1 \dots f_n g_1 \dots g_{n-1})$$

and the R_{new}-module

$$W = R_{\text{new}} \oplus (u, f_1) \oplus (u, f_1 g_1) \oplus (u, f_1 g_1 f_2) \oplus (u, f_1 g_1 f_2 g_2) \oplus \dots \oplus (u, f_1 g_1 f_2 \dots g_{n-1}).$$

By 5.17 W is a MM R_{new}-module. By 5.21 there are no loops in the quiver of $\operatorname{End}_{R_{\text{new}}}(W)$, hence by 5.23 W is a CT R_{new}-module. This implies that all MM R_{new}-generators are CT (2.11), hence in particular

$$Y := \left(\bigoplus_{i=1}^{n} (u, f_1 \dots f_i) \right) \oplus \left(\bigoplus_{j=1}^{n-1} (u, f_1 \dots f_n g_1 \dots g_j) \right)$$

is a CT R_{new}-module. By 5.11(3) it follows that

$$\left(\bigoplus_{i=1}^{n} \frac{S}{f_1 \dots f_i} \right) \oplus \left(\bigoplus_{j=1}^{n-1} \frac{S}{f_1 \dots f_n g_1 \dots g_j} \right)$$

is a CT$^\flat$ $R^\flat_{\text{new}} := S/f_1 \dots f_n g_1 \dots g_{n-1}$-module. Now repeatedly applying 5.25 to the chain of surjections

$$\frac{S}{f_1 \dots f_n g_1 \dots g_{n-1}} \twoheadrightarrow \frac{S}{f_1 \dots f_n g_1 \dots g_{n-2}} \twoheadrightarrow \dots \twoheadrightarrow \frac{S}{f_1 \dots f_n g_1} \twoheadrightarrow \frac{S}{f_1 \dots f_n} = R^\flat$$

yields that $\oplus_{i=1}^{n} S/f_1 \dots f_i$ is a CT$^\flat$ $R^\flat$-module, as required.

The final statement follows since, as in the proof of (2)⇒(3), if there exists a CT R-module, MM generators are precisely the CT R-modules (2.11). Hence since $\{T^\omega \mid \omega \in \mathfrak{S}_n\}$ give all MM generators (by 5.16), these are all the CT R-modules. It follows from 5.11(3) that $\{S^\omega \mid \omega \in \mathfrak{S}_n\}$ are all the $\mathrm{CT}^\flat$ $R^\flat$-modules. $\square$

6. Mutation

6.1. **Some theory.** An operation called mutation was introduced in [IW10, §6]. Although similar in spirit to the mutation of Fomin–Zelevinsky, this operation is categorical, and is designed to deal with situations (like here) where the quivers have many loops and two-cycles, and the relations are not necessarily given by a potential. For our purposes in future subsections we need slightly more information than is given in [IW10, §6], so we record that necessary information here. The main new result is 6.3.

In this subsection we let R be a complete local normal d-CY^- commutative ring (i.e. a complete local normal d-dimensional commutative Gorenstein ring) with $d \geq 2$, and we let $M := R \oplus (\oplus_{i\in I} M_i)$ be a modifying R-module, where we can (and will) assume that M is basic, i.e. all summands are pairwise non-isomorphic. Necessarily $M \in \mathrm{CM}\, R$. We denote $\mathrm{Hom}_R(-, R) := (-)^* : \mathrm{ref}\, R \to \mathrm{ref}\, R$ to be the duality functor and we set $\Lambda_I := \mathrm{End}_R(M)$.

Choose $\emptyset \neq J \subseteq I$, set $M_J := \oplus_{j\in J} M_j$ and $\frac{M}{M_J} := R \oplus (\oplus_{i\in I\backslash J} M_i)$. For convenience we denote $\mathbb{G} := \mathrm{Hom}_R(\frac{M^*}{M_J^*}, -)$ and $\mathbb{F} := \mathrm{Hom}_R(M, -)$. Let

$$U_J^* \xrightarrow{f} M_J^*$$

denote a minimal right add $\frac{M^*}{M_J^*}$-approximation of M_J^*. Recall that a morphism $a : X \to Y$ is called *right minimal* if any $f \in \mathrm{End}_R(X)$ satisfying $a = fa$ is an automorphism. We denote the kernel by K_J. Since $\frac{M^*}{M_J^*}$ is a generator, necessarily f is surjective. Hence $K_J \in \mathrm{CM}\, R$, so by definition we have a short exact sequence

$$0 \to K_J \to U_J^* \xrightarrow{f} M_J^* \to 0 \tag{6.I}$$

in $\mathrm{CM}\, R$ such that

$$0 \to \mathbb{G}K_J \to \mathbb{G}U_J^* \xrightarrow{\cdot f} \mathbb{G}M_J^* \to 0 \tag{6.J}$$

is exact.

Definition 6.1. [IW10, 6.12] *With notation as above, define the mutation of M at summand(s) M_J to be $\nu_J(M) := \frac{M}{M_J} \oplus K_J^*$.*

Now applying duality $(-)^*$ to (6.I) gives an exact sequence

$$0 \to M_J \xrightarrow{f^*} U_J \to K_J^* \to 0$$

from which applying $\mathbb{F}$ yields an injection

$$0 \to \mathbb{F}M_J \xrightarrow{\mathbb{F}f^*} \mathbb{F}U_J \tag{6.K}$$

Definition 6.2. *We define $Q_J := \mathrm{Cok}(\mathbb{F}f^*)$ and denote $V_J := \mathbb{F}(\frac{M}{M_J}) \oplus Q_J$.*

The aim of this subsection is to prove the following result, which is a simultaneous generalization of [IW10, 6.8] and [IW10, 6.13]. Note that in this paper, when we say tilting module we mean a tilting module of projective dimension less than or equal to one.

Theorem 6.3. *Let R be a complete local normal d-CY^- commutative ring, $M := R \oplus (\oplus_{i\in I} M_i)$ be a basic modifying R-module and choose any $\emptyset \neq J \subseteq I$. Then* $\mathrm{End}_R(M)$ *and* $\mathrm{End}_R(\nu_J(M))$ *are derived equivalent, via the tilting* $\mathrm{End}_R(M)$*-module V_J.*

The proof of 6.3 requires the following technical lemma:

Lemma 6.4. *Suppose that Λ is a symmetric R-algebra and $V \in \mathrm{mod}\,\Lambda$ is a height one projective such that* $\mathrm{End}_\Lambda(V)$ *is a symmetric R-algebra. Then* $\mathrm{End}_\Lambda(V) \cong \mathrm{End}_{\Lambda^{\mathrm{op}}}(V^*)^{\mathrm{op}}$.

Proof. Consider the natural ring homomorphism

$$\operatorname{End}_\Lambda(V) \xrightarrow{\psi:=(-)^*} \operatorname{End}_{\Lambda^{\mathrm{op}}}(V^*)^{\mathrm{op}}.$$

Since V is height one projective Λ-module, via the equivalence

$$V_\mathfrak{p} \in \operatorname{ref}\Lambda_\mathfrak{p} \xrightarrow{(-)^*_\mathfrak{p}} \operatorname{ref}\Lambda_\mathfrak{p}^{\mathrm{op}}$$

we have that ψ is a height one isomorphism. But $\Lambda^{\mathrm{op}} \in \operatorname{ref} R$ and by [IR08, 2.4(1)] $V^* \in \operatorname{ref}\Lambda^{\mathrm{op}}$. Clearly V^* is a height one projective Λ^{op}-module, hence by reflexive equivalence [IR08, 2.4(2)(i)] $\operatorname{End}_{\Lambda^{\mathrm{op}}}(V^*)^{\mathrm{op}}$ is reflexive. Furthermore $\operatorname{End}_\Lambda(V)$ is reflexive by assumption, hence ψ, being a height one isomorphism, is actually an isomorphism. ☐

We can now prove 6.3.

Proof. As before, denote $\Lambda_I := \operatorname{End}_R(M)$. We first show that V_J is a tilting Λ_I-module. This is standard and well-known to experts — we just need to show that $\mathbb{F}f^*$ is a left $\operatorname{add}\mathbb{F}(\frac{M}{M_J})$-approximation, i.e.

$$\operatorname{Hom}_{\Lambda_I}(\mathbb{F}U_J, \mathbb{F}(\tfrac{M}{M_J})) \xrightarrow{(\mathbb{F}f^*)\cdot} \operatorname{Hom}_{\Lambda_I}(\mathbb{F}M_J, \mathbb{F}(\tfrac{M}{M_J})) \to 0$$

is exact. Since M is a generator, $\mathbb{F} : \operatorname{mod} R \to \operatorname{mod}\Lambda_I$ is fully faithful [IW10, 2.5(1)], so it suffices to show that

$$\operatorname{Hom}_R(U_J, \tfrac{M}{M_J}) \xrightarrow{f^*\cdot} \operatorname{Hom}_R(M_J, \tfrac{M}{M_J}) \to 0$$

is exact. By duality this is equivalent to showing that

$$\operatorname{Hom}_R(\tfrac{M^*}{M_J^*}, U_J^*) \xrightarrow{\cdot f} \operatorname{Hom}_R(\tfrac{M^*}{M_J^*}, M_J^*) \to 0$$

is exact. But this is precisely (6.J).

We now show that $\operatorname{End}_{\Lambda_I}(V_J) \cong \operatorname{End}_R(\nu_J(M))$. To see this, note first that Λ_I is d-CY$^-$ by 2.10, so since $\operatorname{End}_{\Lambda_I}(V_J)$ is derived equivalent to Λ_I, it is also d-CY$^-$. Hence by [IR08, 3.8(5)] $\operatorname{End}_{\Lambda_I}(V_J)$ is a symmetric R-algebra. Also, $\Lambda_I = \operatorname{End}_R(M)$ is a symmetric R-algebra, by [IR08, 2.4(3)].

Now the exact sequence

$$0 \to \mathbb{F}M_J \xrightarrow{\mathbb{F}f} \mathbb{F}U_J \to \mathbb{F}(K_J^*) \to \operatorname{Ext}^1_R(M, M_J) \to$$

induces a natural Λ_I-module homomorphism $Q_J \xrightarrow{h} \mathbb{F}(K_J^*)$. Since R is normal

$$\operatorname{Ext}^1_R(M, M_J)_\mathfrak{p} = \operatorname{Ext}^1_{R_\mathfrak{p}}(M_\mathfrak{p}, (M_J)_\mathfrak{p}) = 0$$

for all primes $\mathfrak{p} \in \operatorname{Spec} R$ of height one, and so

$$(Q_J)_\mathfrak{p} \cong (\mathbb{F}(K_J^*))_\mathfrak{p} \cong \operatorname{Hom}_{R_\mathfrak{p}}(M_\mathfrak{p}, (K_J^*)_\mathfrak{p})$$

for these primes. But $(K_J^*)_\mathfrak{p}$ is a free $R_\mathfrak{p}$-module and so since M is a generator it follows that Q_J, hence V_J, is a height one projective Λ_I-module. Thus by 6.4

$$\operatorname{End}_{\Lambda_I}(V_J) \cong \operatorname{End}_{\Lambda_I^{\mathrm{op}}}(V_J^*)^{\mathrm{op}}.$$

Now since h is a height one isomorphism, it follows that h^* is a height one isomorphism. But h^* is a morphism between reflexive modules, so h^* must be an isomorphism, hence

$$V_J^* = (\mathbb{F}(\tfrac{M}{M_J}))^* \oplus Q_J^* \cong (\mathbb{F}(\tfrac{M}{M_J}))^* \oplus (\mathbb{F}(K_J^*))^* = (\mathbb{F}(\tfrac{M}{M_J} \oplus K_J^*))^*.$$

Consequently

$$\operatorname{End}_{\Lambda_I}(V_J) \cong \operatorname{End}_{\Lambda_I^{\mathrm{op}}}((\mathbb{F}(\tfrac{M}{M_J} \oplus K_J^*))^*)^{\mathrm{op}} \cong \operatorname{End}_{\Lambda_I}(\mathbb{F}(\tfrac{M}{M_J} \oplus K_J^*))$$

since

$$(\mathbb{F}(\tfrac{M}{M_J} \oplus K_J^*))^* \in \operatorname{ref}\Lambda_I^{\mathrm{op}} \xrightarrow{(-)^*} \operatorname{ref}\Lambda_I$$

is an equivalence. This then yeilds

$$\operatorname{End}_{\Lambda_I}(V_J) \cong \operatorname{End}_{\Lambda_I}(\mathbb{F}(\tfrac{M}{M_J} \oplus K_J^*)) \cong \operatorname{End}_R(\tfrac{M}{M_J} \oplus K_J^*) = \operatorname{End}_R(\nu_J(M)),$$

where the second isomorphism is just reflexive equivalence (see e.g. [IR08, 2.4(2)]). ☐

Remark 6.5. Since (by 6.3) V_J is always a tilting module, for every $\emptyset \neq J \subseteq I$ we have an equivalence

$$T_J := \mathbf{R}\mathrm{Hom}(V_J,-) : \mathrm{D}^{\mathrm{b}}(\mathrm{mod}\,\mathrm{End}_R(M)) \to \mathrm{D}^{\mathrm{b}}(\mathrm{mod}\,\mathrm{End}_R(\nu_J(M))).$$

Note that this functor is never the identity, even although $\nu_J(M) = M$ can happen (see next subsection).

Corollary 6.6. *Let R be a complete local normal d-CY$^-$ commutative ring, $M := R \oplus (\oplus_{i\in I} M_i)$ be a basic modifying R-module and choose any $\emptyset \neq J \subseteq I$. Then $\nu_J(M)$ is a modifying R-module.*

Proof. The proof is identical to the proof of 2.19(1). ☐

Hence the mutation operator ν_J acts on the set of modifying modules. In dimension three, we can say more.

Corollary 6.7. *With notation as in 6.6,*
(1) *If M is maximal modifying, so is $\nu_J(M)$.*
(2) *If M is CT, so is $\nu_J(M)$.*

Proof. Is immediate from 2.19. ☐

Hence in dimension three, the mutation operator ν_J acts on the set of maximal modifying (respectively CT) modules.

6.2. **Mutation of modifying generators on cA_n singularities.** As previously, we let $R := \Bbbk[[u,v,x,y]]/(uv-f)$, where $f = f_1 \dots f_n$ is a factorization into (not necessarily distinct) prime elements f_i.

Even although we do not know if the derived equivalence in §4.1 holds for arbitrary fields, given some (possibly non-maximal) flag $\mathcal{F} = (I_1,\dots,I_m)$ in the set $\{1,2,\dots,n\}$, it is useful combinatorially (as in 4.6) to assign to $\mathcal{F}$ the following picture:

$\mathcal{P}(\mathcal{F}) :=$ 

where $g_j := \frac{f_{I_j}}{f_{I_{j-1}}}$ for all $1 \leq j \leq m+1$, where by convention $I_0 := \emptyset$ and $I_{m+1} := \{1,2,\dots,n\}$.

There are m curves. If we denote $M_0 := R$ and $M_j := (u, f_{I_j}) = (u, \prod_{i=1}^{j} g_i)$, and set $T^{\mathcal{F}} := \oplus_{j=0}^{m} M_j$, then $T^{\mathcal{F}}$ is the modifying R-module corresponding to $\mathcal{F}$. The correpondence between non-free summands of $T^{\mathcal{F}}$ and curves of $\mathcal{P}(\mathcal{F})$ is as follows:



This gives us a combinatorial model to visualize mutation.

Example 6.8. Consider $f = f_1f_2f_3f_4f_5f_6$ with flag $\mathcal{F} = (\{2,3\} \subsetneq \{2,3,1\})$. Then $\mathcal{F}$ corresponds to



The correponding $T^{\mathcal{F}}$ is $(u, f_2f_3) \oplus (u, f_1f_2f_3) \oplus R$.

We are interested in mutations of non-free summands of $T^{\mathcal{F}}$, so since above such summands correspond to subsets of the curves, pick an arbitrary $\emptyset \neq J \subseteq \{1,\dots,m\}$. Now write J as a disjoint union of connected components:

Definition 6.9. *A connected component of J is a collection of consecutive numbers from i_1 to i_2 inside $\{1,\dots,n\}$, each of which belongs to J, such that $i_1 - 1 \notin J$ and $i_2 + 1 \notin J$. We write $J = \coprod_{j=1}^{t} J_j$ as a disjoint union of connected components.*

Geometrically, if say $m = 6$ and $J := \{2,3,5\}$ then we are simply bunching the curves corresponding to J into connected components as in the following picture:



The mutation operation acts on the set of modifying modules, hence on $T^{\mathcal{F}}$, and hence on the set of $\mathcal{P}(\mathcal{F})$. Below (see 6.13) we will justify the following intuitive geometric picture:



where each connected component of J gets reflected. It is clear from this picture (and indeed we prove it in 6.13) that $\nu_J(M) = M$ if and only if J is componentwise symmetric.

Definition 6.10. *For a given flag $\mathcal{F} = (I_1, \ldots, I_m)$, associate the combinatorial picture $\mathcal{P}(\mathcal{F})$ as above. For $\emptyset \neq J \subseteq \{1, \ldots, m\}$, we define the J-reflection of $\mathcal{P}(\mathcal{F})$ as follows: the number of curves remains the same, but the new position of the g's is obtained from the positions in $\mathcal{P}(\mathcal{F})$ by reflecting each connected component of J in the vertical axis.*

We now build up to 6.13. To fix some convenient notation, we write $J = \coprod_{j=1}^{t} J_j$ and denote

$$J_j = \{l_j, l_j + 1, \ldots, u_j - 1, u_j\}$$

to be the connected components of J. The l_j stands for lower bound, whilst u_j stands for upper bound.

Lemma 6.11. *Fix flag $\mathcal{F} = (I_1, \ldots, I_m)$, and choose $\emptyset \neq J \subseteq \{1, \ldots, m\}$. Write $J = \coprod_{j=1}^{t} J_j$ and consider one of the summands M_i of $T^{\mathcal{F}}$ which lies in J. Say M_i lives in the component J_j, then the following sequence is exact*

$$0 \to M_i \xrightarrow{(inc \ \prod_{a=i+1}^{u_j+1} g_a)} M_{l_j-1} \oplus M_{u_j+1} \xrightarrow{\left(\begin{smallmatrix} \prod_{a=i+1}^{u_j+1} g_a \\ -inc \end{smallmatrix}\right)} (u, (\prod_{b=1}^{l_j-1} g_b)(\prod_{a=i+1}^{u_j+1} g_a)) \to 0. \tag{6.L}$$

Proof. Note first the maps go where they should. The right hand map is clearly surjective since the generators of the module $(u, (\prod_{b=1}^{l_j-1} g_b)(\prod_{a=i+1}^{u_j+1} g_b))$ are both in the image of the map. The left hand map is clearly injective. An easy check shows that the above is exact in the middle. ☐

Lemma 6.12. *The dual short exact sequence of (6.L), namely*

$$0 \to (u, (\textstyle\prod_{b=1}^{l_j-1} g_b)(\prod_{a=i+1}^{u_j+1} g_a))^* \to M_{l_j-1}^* \oplus M_{u_j+1}^* \to M_i^* \to 0,$$

is a minimal right add $\frac{M^*}{M_J^*}$*-approximation of M_i^*.*

Proof. This follows immediately from the shape of the quiver of $\operatorname{End}_R(T^{\mathcal{F}})$ in 5.21. ☐

Theorem 6.13. *Fix a flag $\mathcal{F} = (I_1, \ldots, I_m)$, and associate to $\mathcal{F}$ the module $T^{\mathcal{F}}$ and the combinatorial picture $\mathcal{P}(\mathcal{F})$, as before. Choose $\emptyset \neq J \subseteq \{1, \ldots, m\}$, then $\nu_J(T^{\mathcal{F}})$ is the module corresponding to the J-reflection of $\mathcal{P}(\mathcal{F})$.*

Proof. Consider the first connected component J_1. By 6.12 we know that M_{u_1} mutates to $(u, (\prod_{b=1}^{l_1-1} g_b) g_{u_1+1})^{**}$, which is isomorphic to $(u, (\prod_{b=1}^{l_1-1} g_b) g_{u_1+1})$. Similarly, M_{u_1-1} mutates to $(u, (\prod_{b=1}^{l_1-1} g_b) g_{u_1+1} g_{u_1})$. Continuing, we see that M_{u_1-i} mutates to $(u, (\prod_{b=1}^{l_1-1} g_b) g_{u_1+1} \ldots g_{u_1-i+1})$ for all $1 \leq i \leq u_1 - l_1$. Since the combinatorial picture is built by ordering the summands in increasing lengths of products, we see that in the combinatorial picture, the component J_1 has been reflected. The proof that the remaining components are reflected is identical. ☐

Example 6.14. As in 6.8, consider $f = f_1 f_2 f_3 f_4 f_5 f_6$ with flag $\mathcal{F} = (\{2,3\} \subsetneq \{2,3,1\})$. Then $T^{\mathcal{F}} = R \oplus (u, f_2 f_3) \oplus (u, f_1 f_2 f_3)$, which pictorially is



Pick summand $(u, f_1 f_2 f_3)$, then the mutation is given by



and so $\nu(T^{\mathcal{F}}) = R \oplus (u, f_2 f_3) \oplus (u, f_2 f_3 f_4 f_5 f_6)$.

6.3. **Geometric corollaries.** We set $k = \mathbb{K}$. Given 4.2, 6.3 and 6.13, we can now produce derived equivalences between partial crepant resolutions of $\operatorname{Spec} R$. First, note that even if both contain the same number of curves above the origin (and hence their K-theory match), two partial resolutions need not be derived equivalent:

Example 6.15. In the case $f = xxy$, the crepant partial resolutions with one curve are



$X^{\{1\}}$



$X^{\{1,3\}}$



$X^{\{2\}}$



$X^{\{1,2\}}$

The two spaces $X^{\{1\}}$ and $X^{\{1,3\}}$ are derived equivalent via mutation, and the two spaces $X^{\{2\}}$ and $X^{\{1,2\}}$ are also derived equivalent via mutation. However they are not all derived equivalent, since if $\mathrm{D}^{\mathrm{b}}(\operatorname{coh} X^{\{1\}}) \approx \mathrm{D}^{\mathrm{b}}(\operatorname{coh} X^{\{2\}})$ then $\mathrm{D}_{\mathsf{sg}}(X^{\{1\}}) \approx \mathrm{D}_{\mathsf{sg}}(X^{\{2\}})$, which by 3.7 gives $\underline{\mathrm{CM}}\,\mathbb{K}[[u,v,x,y]]/(uv - x^2) \approx \underline{\mathrm{CM}}\,\mathbb{K}[[u,v,x,y]]/(uv - xy)$. But this is impossible, since (for example) the left hand side has infinite dimensional Hom-spaces, whereas in the right hand side all Hom spaces are finite dimensional.

The following is immediate from the theory of mutation:

Corollary 6.16. *In the case $k = \mathbb{K}$, suppose that $\mathcal{F} = (I_1, \ldots, I_m)$ and $\mathcal{G} = (J_1, \ldots, J_n)$ are flags. Then $X^{\mathcal{F}}$ and $X^{\mathcal{G}}$ are derived equivalent if $n = m$ and the singularities of $X^{\mathcal{F}}$ can be permuted to the singularities of $X^{\mathcal{G}}$.*

Proof. By 4.2 $\mathrm{D}^{\mathrm{b}}(X^{\mathcal{F}}) = \mathrm{D}^{\mathrm{b}}(\operatorname{End}_R(T^{\mathcal{F}}))$ and $\mathrm{D}^{\mathrm{b}}(X^{\mathcal{G}}) = \mathrm{D}^{\mathrm{b}}(\operatorname{End}_R(T^{\mathcal{G}}))$, hence the result follows if we establish that $\operatorname{End}_R(T^{\mathcal{F}})$ and $\operatorname{End}_R(T^{\mathcal{G}})$ are derived equivalent. Now mutation always gives derived equivalences (6.3), and mutation corresponds to permuting the order of the singularities (6.13). Hence if the singularities of $X^{\mathcal{F}}$ can be permuted to the singularities of $X^{\mathcal{G}}$, certainly there is a finite number of mutations which transforms $\mathcal{P}(\mathcal{F})$ into $\mathcal{P}(\mathcal{G})$, hence $\operatorname{End}_R(T^{\mathcal{F}})$ and $\operatorname{End}_R(T^{\mathcal{G}})$ are derived equivalent. $\square$

6.4. **The mutation graph for maximal modifying modules.** By 6.15 and 6.16, for general partial resolutions it is clear that very rarely will $T^{\mathcal{F}_1}$ and $T^{\mathcal{F}_2}$ be linked by mutations.

However, if both $T^{\mathcal{F}_1}$ and $T^{\mathcal{F}_2}$ have the maximal number of summands (i.e. $\mathcal{P}(\mathcal{F}_1)$ and $\mathcal{P}(\mathcal{F}_2)$ have the maximal number of curves), the homological algebra is much better behaved. We now prove, when $k = \Bbbk$, that the mutation graph for MM generators is connected. Combining this with 6.16 in the case $k = \mathbb{C}$ proves 1.12 in the introduction, namely that all $\mathbb{Q}$-factorial terminalizations of $\operatorname{Spec} R$ are derived equivalent.

Recall that by the *mutation graph* of MM generators we mean the graph in which the vertices are the (isomorphism classes of) basic MM generators, where we draw an edge between two vertices M and N if there exists a mutation ν_J for which $\nu_J(M) \cong N$.

Corollary 6.17. *Let $R := \Bbbk[[u,v,x,y]]/(uv - f)$, where $f \in (x,y) \subseteq \Bbbk[[x,y]]$. Then the mutation graph of MM generators is connected.*

Proof. As before, write $f = f_1 \ldots f_n$ as a factorization into (not necessarily distinct) prime elements f_i. Suppose M and N are MM generators, then by the above we can view M and N combinatorially as





where $\{j_1, \dots, j_n\} = \{k_1, \dots, k_n\} = \{1, \dots, n\}$. Now mutation corresponds to permutations, and it is clear that we can transform $\mathcal{P}(M)$ into $\mathcal{P}(N)$ by a finite sequence of permutations. It follows that the mutation graph is connected. $\square$

Example 6.18. Let $f = f_1^2 f_2^3 = f_1 f_1 f_2 f_2 f_2$. Then the mutation graph for maximal modifying generators is the following


$f_1\ f_1\ f_2\ f_2\ f_2$
$f_2\ f_1\ f_1\ f_2\ f_2$
$f_2\ f_2\ f_1\ f_1\ f_2$
$f_2\ f_2\ f_2\ f_1\ f_1$
$f_1\ f_2\ f_1\ f_2\ f_2$
$f_2\ f_1\ f_2\ f_1\ f_2$
$f_2\ f_2\ f_1\ f_2\ f_1$
$f_1\ f_2\ f_2\ f_1\ f_2$
$f_2\ f_1\ f_2\ f_2\ f_1$
$f_1\ f_2\ f_2\ f_2\ f_1$

where for clarity we have illustrated only those that are connected via a mutation by a single summand. If we include mutations by more that one summand, there are many more connecting lines.

Remark 6.19. Geometers will recognize the above picture, since when $k = \mathbb{C}$ it corresponds exactly to the flops of curves on the $\mathbb{Q}$-factorial terminalizations of $\operatorname{Spec} R$. We will return to this in future.

References

[Aus84] M. Auslander, *Isolated singularities and existence of almost split sequences.* Representation theory, II (Ottawa, Ont., 1984), 194–242, Lecture Notes in Math., 1178, Springer, Berlin, 1986.

[AG60] M. Auslander and O. Goldman, *Maximal orders*, Trans. Amer. Math. Soc. **97** (1960), 1–24.

[Bri02] T. Bridgeland, *Flops and derived categories.* Invent. Math. **147** (2002), no. 3, 613–632

[BKR] T. Bridgeland, A. King and M. Reid, *The McKay correspondence as an equivalence of derived categories*, J. Amer. Math. Soc. **14** (2001), no. 3, 535–554.

[BH] W. Bruns and J. Herzog, *Cohen-Macaulay rings. Rev. ed.*, Cambridge Studies in Advanced Mathematics. **39** xiv+453 pp.

[Bu86] R. -O. Buchweitz, *Maximal Cohen–Macaulay modules and Tate–cohomology over Gorenstein rings*, preprint, 1986.

[BIKR] I. Burban, O. Iyama, B. Keller and I. Reiten, *Cluster tilting for one-dimensional hypersurface singularities*, Adv. Math. **217** (2008), no. 6, 2443–2484.

[BK11] I. Burban and M. Kalck, *Singularity category of a non-commutative resolution of singularities*, arXiv:1103.3936.

[C02] J-C. Chen, *Flops and equivalences of derived categories for threefolds with only terminal Gorenstein singularities*, J. Differential Geom. **61** (2002), no. 2, 227–261.

[C10] X-W. Chen, *Unifying two results of Orlov on singularity categories*, Abh. Math. Semin. Univ. Hambg. **80** (2010), no. 2, 207–212.

[CYZ] X-W. Chen, Y. Ye and P. Zhang, *Algebras of derived dimension zero*, Comm. Algebra **36** (2008), no. 1, 1–10.

[Dao1] H. Dao, *Remarks on non-commutative crepant resolutions of complete intersections*, arXiv:0911.4541.

[Dao2] H. Dao, *Decency and rigidity over hypersurfaces*, arXiv:AC/0611568v1.

[Dao3] H. Dao, *Picard groups of punctured spectra of dimension three local hypersurfaces are torsion-free*, arXiv:1004.0471.

[DH10] H. Dao and C. Huneke, *Vanishing of Ext, Cluster Tilting and Finite Global Dimension of Endomorphism Rings*, arXiv:1005.5359 (2010).

[F77] H–B. Foxby, *Isomorphisms between complexes with applications to the homological theory of modules*, Mathematica Scandinavica, **40**, (1977) no.1, 5–19.

[GR71] L. Gruson and M. Raynaud, *Critères de platitude et de projectivité. Techniques de 'platification' d'un module*, Invent. Math. **13**, (1971) 1–89.

[H77] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, No. 52. Springer-Verlag, 1977. xvi+496 pp.

[IR08] O. Iyama and I. Reiten, *Fomin-Zelevinsky mutation and tilting modules over Calabi-Yau algebras*, Amer. J. Math. **130** (2008), no. 4, 1087–1149.

[IW08] O. Iyama and M. Wemyss, *The classification of special Cohen Macaulay modules*, Math. Z. **265** (2010), no.1, 41–83.

[IW10] O. Iyama and M. Wemyss, *Auslander-Reiten Duality and Maximal Modifications for Non-isolated Singularities*, arXiv:1007.1296.

[J70] C. U. Jensen, *On the vanishing of* $\varprojlim^{(i)}$ J. Alg. **15**, (1970) 151–166.

[Jo75] P. Jothilingam, *A note on grade*, Nagoya Math. J. **59** (1975), 149 –152.

[K88] Y. Kawamata, *Crepant blowing-up of 3-dimensional canonical singularities and its application to degenerations of surfaces*, Ann. of Math. (2) **127** (1988), no. 1, 93–163.

[KMV] B. Keller, D. Murfet, M. Van den Bergh, *On two examples by Iyama and Yoshino*, arXiv:0803:0720.

[KM98] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Math., **134**, Cambridge Univ. Press, Cambridge, 1998.

[N01] A. Neeman, *Triangulated categories*, Annals of Mathematics Studies, vol. 148, Princeton University Press, Princeton, NJ, 2001.

[O76] A. Ooishi, *Matlis duality and the width of a module*, Hiroshima Math. J. **6** (1976), 573–587.

[Or03] D. Orlov, *Triangulated categories of singularities and D-branes in Landau-Ginzburg models*, arXiv:AG/0302304.

[Or05] D. Orlov, *Triangulated categories of singularities and equivalences between Landau-Ginzburg Models*, arXiv:AG/0503630.

[Or09] D. Orlov, *Formal completions and idempotent completions of triangulated categories of singularities*, arXiv:0901.1859.

[R83] M. Reid, *Minimal models of canonical 3-folds*, Algebraic varieties and analytic varieties (Tokyo, 1981), 131–180, Adv. Stud. Pure Math., 1, North-Holland, Amsterdam, 1983.

[R87] M. Reid, *Young person's guide to canonical singularities*, Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985), 345–414, Proc. Sympos. Pure Math., 46, Part 1, Amer. Math. Soc., Providence, RI, 1987.

[R89] J. Rickard, *Morita theory for derived categories*, J. London Math. Soc. (2) **39** (1989), 436–456.

[R98] P. Roberts, *Multiplicities and Chern classes in local algebra*, Cambridge Tracts in Mathematics, 133. Cambridge University Press, Cambridge, 1998. xii+303 pp.

[R08] R. Rouquier, *Dimensions of triangulated categories*, J. K-Theory **1** (2008), no. 2, 193–256.

[S89] O. Solberg, *Hypersurface singularities of finite Cohen-Macaulay type*. Proc. London Math. Soc. (3) **58** (1989), no. 2, 258–280.

[V04a] M. Van den Bergh, *Three-dimensional flops and noncommutative rings*, Duke Math. J. **122** (2004), no. 3, 423–455.

[V04b] ———, *Non-commutative crepant resolutions*, The legacy of Niels Henrik Abel, 749–770, Springer, Berlin, 2004.

[W10] M. Wemyss, *The* $\mathrm{GL}(2,\mathbb{C})$ *McKay correspondence*, Math. Ann. **350**, no. 3, (2011), 631–659.

[Yos90] Y. Yoshino, *Cohen-Macaulay modules over Cohen-Macaulay rings*, London Mathematical Society Lecture Note Series, 146. Cambridge University Press, Cambridge, 1990.

Osamu Iyama, Graduate School of Mathematics, Nagoya University, Chikusa-ku, Nagoya, 464-8602, Japan

E-mail address: `iyama@math.nagoya-u.ac.jp`

Michael Wemyss, School of Mathematics, James Clerk Maxwell Building, The King's Buildings, Mayfield Road, Edinburgh, EH9 3JZ, UK.

E-mail address: `wemyss.m@googlemail.com`